

11/22/99

Please type a plus sign (+) inside this box → ☐

Approved for use through 09/30/2000. OMB 0951-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))

Attorney Docket No. 6356.US.P3
First Inventor or Application Identifier J. Henkin, et al.
Title PEPTIDE ANTIANGIOGENIC DRUGS
Express Mail Label No. EL384166875US

APPLICATION ELEMENTS
See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

1. ☒ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original and a duplicate for fee processing)
2. ☒ Specification [Total Pages 229]
(preferred arrangement set forth below)
 - Descriptive title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings (if filed)
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
3. ☐ Drawing(s) (35 U.S.C. 113) [Total Sheets ☐
4. Oath or Declaration [Total Pages ☐- a. ☐ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 C.F.R. § 1.63(d))
(for continuation/divisional with Box 16 completed)
 - i. ☐ **DELETION OF INVENTOR(S)**
Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).

5. ☐ Microfiche Computer Program (Appendix)
6. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy (identical to computer copy)
 - c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

7. ☐ Assignment Papers (cover sheet & document(s))
8. ☐ 37 C.F.R. §3.73(b) Statement ☐ Power of Attorney (when there is an assignee)
9. ☐ English Translation Document (if applicable)
10. ☒ Information Disclosure Statement (IDS)/PTO-1449 ☒ Copies of IDS Citations
11. ☐ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
13. ☐ Small Entity ☐ Statement filed in prior application (PTO/SB-92-12) Status still proper and desired
14. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
15. ☒ Other: Declaration/Power of Attorney (Unexecuted)

* NOTE FOR ITEMS 1 & 13 IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

☐ Continuation ☐ Divisional ☒ Continuation-in-part (CIP) of prior application No: 09/316,888
Prior application information: Examiner F. Moezie Group / Art Unit: 1646

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

17. CORRESPONDENCE ADDRESS

☐ Customer Number or Bar Code Label (Insert Customer No. or Attach bar code label here) or ☒ Correspondence address below

Name	<u>Steven F. Weinstock</u>				
	<u>Abbott Laboratories</u>				
Address	<u>Dept. 377 Bldg. AP6D-2</u>				
	<u>100 Abbott Park Road</u>				
City	<u>Abbott Park</u>	State	<u>IL</u>	Zip Code	<u>60064-6050</u>
Country	<u>United States of America</u>	Telephone	<u>(847) 937-4558</u>	Fax	<u>(847) 938-2623</u>

Name (Print/Type) Portia Chen Registration No. (Attorney/Agent) 44,075
Signature [Signature] Date 11/22/99

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

PEPTIDE ANTIANGIOGENIC DRUGS

This application claims priority from the U.S. application Ser. No. 09/316,888, filed May 21, 1999, which claims priority from U.S. application Ser No. 60/086,536, filed May 22, 1998, and U.S. application Ser. No. 60/126,546, filed March 26, 1999.

Technical Field

The invention relates to novel compounds having activity useful for treating conditions which arise or are exacerbated by angiogenesis, pharmaceutical compositions comprising these compounds, a method of treating using said compounds, and a method of inhibiting angiogenesis.

Background of the Invention

Angiogenesis is the fundamental process by which new blood vessels are formed and is essential to a variety of normal body activities (such as reproduction, development and wound repair). Although the process is not completely understood, it is believed to involve a complex interplay of molecules which both stimulate and inhibit the growth of endothelial cells, the primary cells of the capillary blood vessels. Under normal conditions, these molecules appear to maintain the microvasculature in a quiescent state (i.e. one of no capillary growth) for prolonged periods which may last for as long as weeks or in some cases, decades. When necessary however (such as during wound repair), these same cells can undergo rapid proliferation and turnover within a five day period. (Folkman, J. and Shing, Y., *The Journal of Biological Chemistry*, **267**(16): 10931-10934, and Folkman, J. and Klagsbrun, M., *Science*, **235**: 442-447 (1987)).


Although angiogenesis is a highly regulated process under normal conditions, many diseases (characterized as "angiogenic diseases") are driven by persistent unregulated angiogenesis. Otherwise stated, unregulated angiogenesis may either cause a particular disease directly or exacerbate an existing pathological condition. For example, ocular neovascularization has been implicated as the most common cause of blindness. In certain existing conditions such as arthritis, newly formed capillary blood vessels invade the joints and destroy cartilage. In diabetes, new capillaries formed in the retina invade the vitreous, bleed, and cause blindness. Growth and metastasis of solid tumors are also angiogenesis-dependent (Folkman, J., *Cancer Research*, **46**: 467-473 (1986), Folkman, J., *Journal of the National Cancer Institute*, **82**: 4-6 (1989)). It has been shown for example that tumors which enlarge to greater than 2 mm, must obtain their own blood supply and do so by inducing the growth of new capillary blood vessels. Once these new blood vessels become embedded in the tumor, they provide a means for tumor cells to enter the

circulation and metastasize to distant sites, such as liver, lung or bone (Weidner, N., et al., *The New England Journal of Medicine*, **324**(1): 1-8 (1991)).

Although several angiogenesis inhibitors are currently under development for use in treating angiogenic diseases (Gasparini, G. and Harris, A.L., *J Clin Oncol* **13**(3): 765-782, (1995)), there are disadvantages associated with several of these compounds. For example, suramin is a potent angiogenesis inhibitor, but causes (at doses required to reach antitumor activity) severe systemic toxicity in humans. Other compounds, such as retinoids, interferons and antiestrogens are safe for human use but have only a weak anti-angiogenic effect.

Summary of the Invention

In one aspect, the present invention provides a compound the formula of:



or a pharmaceutically acceptable salt, ester, solvate or prodrug thereof, wherein:

A_0 is hydrogen or an acyl group selected from:

- (1) $R-(CH_2)_n-C(O)-$; wherein n is an integer from 0 to 8 and R is selected from hydroxyl; methyl; N-acetylamino; methoxy; carboxyl; cyclohexyl optionally containing a one or two double bonds and optionally substituted with one to three hydroxyl groups; and a 5- or 6-membered ring aromatic or nonaromatic ring optionally containing one or two heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the ring is optionally substituted with a moiety selected from alkyl, alkoxy, and halogen; and
- (2) $R^1-CH_2CH_2-(OCH_2CH_2O)_p-CH_2-C(O)-$; wherein R^1 is selected from hydrogen, alkyl and N-acetylamino, and p is an integer from 1 to 8;

A_1 is an amino acyl residue selected from:

- (1) alanyl,
- (2) asparaginylyl,
- (3) citrullyl,
- (4) glutaminylyl,

- 5
- (5) glutamyl,
 - (6) N-ethylglycyl,
 - (7) methionyl,
 - (8) N-methylalanyl,
 - (9) prolyl,
 - (10) pyro-glutamyl,
 - (11) sarcosyl,
 - (12) seryl,
 - (13) threonyl,
- 10
- (14) $-\text{HN}-(\text{CH}_2)_q-\text{C}(\text{O})-$, wherein q is 1 to 8, and
 - (15) $-\text{HN}-\text{CH}_2\text{CH}_2-(\text{OCH}_2\text{CH}_2\text{O})_r-\text{CH}_2-\text{C}(\text{O})-$, wherein r is 1 to 8;

A₂ is an amino acyl residue selected from:

- 15
- (1) alanyl,
 - (2) asparaginylyl,
 - (3) aspartyl,
 - (4) glutaminyl,
 - (5) glutamyl,
 - (6) leucyl,
 - (7) methionyl,
- 20
- (8) phenylalanyl,
 - (9) prolyl,
 - (10) seryl,
 - (11) $-\text{HN}-(\text{CH}_2)_q-\text{C}(\text{O})-$, wherein q is 1 to 8, and
 - (12) $-\text{HN}-\text{CH}_2\text{CH}_2-(\text{OCH}_2\text{CH}_2\text{O})_r-\text{CH}_2-\text{C}(\text{O})-$, wherein r is 1 to 8;

A₃ is an amino acyl residue selected from:

- 25
- (1) alanyl,
 - (2) asparaginylyl,
 - (3) citrullyl,
 - (4) cyclohexylalanyl,
- 30
- (5) cyclohexylglycyl,
 - (6) glutaminyl,
 - (7) glutamyl,











- (8) glycyl,
- (9) isoleucyl,
- (10) leucyl,
- (11) methionyl,
- (12) norvalyl,
- (13) phenylalanyl,
- (14) seryl,
- (15) *t*-butylglycyl,
- (16) threonyl,
- (17) valyl,
- (18) penicillaminyl, and
- (19) cystyl;

A₄ is an amino acyl residue of L or D configuration selected from:

- (1) allo-isoleucyl,
- (2) glycyl,
- (3) isoleucyl,
- (4) prolyl,
- (5) dehydroleucyl,
- (6) D-alanyl,
- (7) D-3-(naphth-1-yl)alanyl,
- (8) D-3-(naphth-2-yl)alanyl,
- (9) D-(3-pyridyl)-alanyl,
- (10) D-2-aminobutyryl,
- (11) D-allo-isoleucyl,
- (12) D-allo-threonyl;
- (13) D-allylglycyl,
- (14) D-asparaginyI,
- (15) D-aspartyl,
- (16) D-benzothienyl,
- (17) D-3-(4,4'-biphenyl)alanyl,
- (18) D-chlorophenylalanyl,
- (19) D-3-(3-trifluoromethylphenyl)alanyl,

	(20)	D-3-(3-cyanophenyl)alanyl,
	(21)	D-3-(3,4-difluorophenyl)alanyl,
	(22)	D-citrullyl,
5	(23)	D-cyclohexylalanyl,
	(24)	D-cyclohexylglycyl,
	(25)	D-cystyl,
	(26)	D-cystyl(<i>S</i> - <i>t</i> -butyl),
	(27)	D-glutaminyI,
	(28)	D-glutamyl,
10	(29)	D-histidyl,
	(30)	D-homoisoleucyl,
	(31)	D-homophenylalanyl,
	(32)	D-homoseryl,
15	(33)	D-isoleucyl,
	(34)	D-leucyl,
	(35)	D-lysyl(N-epsilon-nicotinyl),
	(36)	D-lysyl,
	(37)	D-methionyl,
20	(38)	D-neopentylglycyl,
	(39)	D-norleucyl,
	(40)	D-norvalyl,
	(41)	D-ornithyl,
	(42)	D-penicillaminyI,
	(43)	D-penicillaminyI(acetamidomethyl),
25	(44)	D-penicillaminyI(<i>S</i> -benzyl),
	(45)	D-phenylalanyl,
	(46)	D-3-(4-aminophenyl)alanyl,
	(47)	D-3-(4-methylphenyl)alanyl,
	(48)	D-3-(4-nitrophenyl)alanyl,
30	(49)	D-3-(3,4-dimethoxyphenyl)alanyl,
	(50)	D-3-(3,4,5-trifluorophenyl)alanyl,
	(51)	D-prolyl,

- 5 (52) D-seryl,
 (53) D-seryl(*O*-benzyl),
 (54) D-*t*-butylglycyl,
 (55) D-thienylalanyl,
 (56) D-threonyl,
 (57) D-threonyl(*O*-benzyl),
 (58) D-tryptyl,
 (59) D-tyrosyl(*O*-benzyl),
 (60) D-tyrosyl(*O*-ethyl),
 10 (61) D-tyrosyl, and
 (62) D-valyl;

A₅ is an amino acyl residue of L or D configuration selected from:

- 15 (1) alanyl,
 (2) (3-pyridyl)alanyl,
 (3) 3-(naphth-1-yl)alanyl,
 (4) 3-(naphth-2-yl)alanyl,
 (5) allo-threonyl,
 (6) allylglycyl,
 (7) glutaminyl,
 20 (8) glycyl,
 (9) histidyl,
 (10) homoseryl,
 (11) isoleucyl,
 (12) lysyl(N-epsilon-acetyl),
 25 (13) methionyl,
 (14) norvalyl,
 (15) octylglycyl,
 (16) ornithyl,
 (17) 3-(4-hydromethylphenyl)alanyl,
 30 (18) prolyl,
 (19) seryl,
 (20) threonyl,

- 5
- (21) tryptyl,
 - (22) tyrosyl,
 - (23) D-allo-threonyl,
 - (24) D-homoseryl,
 - (25) D-seryl,
 - (26) D-threonyl,
 - (27) penicillaminy, and
 - (28) cystyl;

A₆ is an amino acyl residue of L or D configuration selected from:

- 10
- (1) alanyl,
 - (2) 3-(naphth-1-yl)alanyl,
 - (3) 3-(naphth-2-yl)alanyl,
 - (4) (3-pyridyl)alanyl,
 - (5) 2-aminobutyryl,
- 15
- (6) allylglycyl,
 - (7) arginyl,
 - (8) asparaginy, and
 - (9) aspartyl,
 - (10) citrully,
- 20
- (11) cyclohexylalanyl,
 - (12) glutaminy,
 - (13) glutamyl,
 - (14) glycyl,
 - (15) histidyl,
- 25
- (16) homoalanyl,
 - (17) homoleucyl,
 - (18) homoseryl,
 - (19) isoleucyl,
 - (20) leucyl,
- 30
- (21) lysyl(N-epsilon-acetyl),
 - (22) lysyl(N-epsilon-isopropyl),
 - (23) methionyl(sulfone),

- 5 (24) methionyl(sulfoxide),
 (25) methionyl,
 (26) norleucyl,
 (27) norvalyl,
 (28) octylglycyl,
 (29) phenylalanyl,
 (30) 3-(4-carboxyamidophenyl)alanyl,
 (31) propargylglycyl,
 (32) seryl,
 10 (33) threonyl,
 (34) tryptyl,
 (35) tyrosyl,
 (36) valyl,
 15 (37) D-3-(naphth-1-yl)alanyl,
 (38) D-3-(naphth-2-yl)alanyl,
 (39) D-glutaminyI,
 (40) D-homoseryl,
 (41) D-leucyl,
 (42) D-norvalyl,
 20 (43) D-seryl,
 (44) penicillaminyI, and
 (45) cystyl;

A₇ is an amino acyl residue of L or D configuration selected from:

- 25 (1) alanyl,
 (2) allylglycyl,
 (3) aspartyl,
 (4) citrullyl,
 (5) cyclohexylglycyl,
 (6) glutamyl,
 30 (7) glycyl,
 (8) homoseryl,
 (9) isoleucyl,

- 5
- (10) allo-isoleucyl,
 - (11) leucyl,
 - (12) lysyl(N-epsilon-acetyl),
 - (13) methionyl,
 - (14) 3-(naphth-1-yl)alanyl,
 - (15) 3-(naphth-2-yl)alanyl,
 - (16) norvalyl,
 - (17) phenylalanyl,
 - (18) prolyl,

10

 - (19) seryl,
 - (20) *t*-butylglycyl,
 - (21) tryptyl,
 - (22) tyrosyl,
 - (23) valyl,

15

 - (24) D-allo-isoleucyl,
 - (25) D-isoleucyl,
 - (26) penicillaminyl, and
 - (27) cystyl;

A₈ is an amino acyl residue selected from:

- 20
- (1) 2-amino-4-[(2-amino-pyrimidinyl)]butanoyl,
 - (2) alanyl(3-guanidino),
 - (3) alanyl[3-pyrrolidinyl(2-N-amidino)],
 - (4) alanyl[4-piperidinyl(N-amidino)],
 - (5) arginyl,

25

 - (6) arginyl(N^GN^Gdiethyl),
 - (7) citrullyl,
 - (8) 3-(cyclohexyl)alanyl(4-Nⁱ-isopropyl),
 - (9) glycyl[4-piperidinyl(N-amidino)],
 - (10) histidyl,

30

 - (11) homoarginyl,
 - (12) lysyl,
 - (13) lysyl(N-epsilon-isopropyl),

- (14) lysyl(N-epsilon-nicotinyl),
- (15) norarginyl,
- (16) ornithyl(N-delta-isopropyl),
- (17) ornithyl(N-delta-nicotinyl),
- (18) ornithyl[N-delta-(2-imidazolynyl)],
- (19) [4-amino(N-isopropyl)methyl]phenyl]alanyl,
- (20) 3-(4-guanidinophenyl)alanyl, and
- (21) 3-(4-amino-N-isopropylphenyl)alanyl;

A₉ is an amino acyl residue of L or D configuration selected from:


- (1) 2-amino-buteryl,
- (2) 2-amino-isobutyryl,
- (3) homopropyl,
- (4) hydroxypropyl,
- (5) isoleucyl,
- (6) leucyl,
- (7) phenylalanyl,
- (8) prolyl,
- (9) seryl,
- (10) *t*-butylglycyl,
- (11) 1,2,3,4-tetrahydroisoquinoline-3-carbonyl,
- (12) threonyl,
- (13) valyl,
- (14) D-alanyl, and
- (15) D-prolyl; and

A₁₀ is a hydroxyl group or an amino acid amide is selected from:

- (1) azaglycyclamide,
- (2) D-alanylamide,
- (3) D-alanylethylamide,
- (4) glycylamide,
- (5) glycylethylamide,
- (6) sarcosylamide,
- (7) serylamide,

(8) D-serylamide,

(9) a group represented by the formula



(9) a group represented by the formula $-\text{NH}-\text{R}^4$;

wherein:

s is an integer selected from 0 to 8,

R^2 is selected from hydrogen, alkyl, and a 5- to 6-membered cycloalkyl ring;

R^3 is selected from hydrogen, hydroxy, alkyl, phenyl, alkoxy, and a 5- to 6-membered ring optionally containing from one to two heteroatoms selected from oxygen, nitrogen, and sulfur, provided that s is not zero when R^3 is hydroxy or alkoxy; and

R^4 is selected from hydrogen, hydroxy, and a 5- to 6-membered cycloalkyl ring.

In another aspect, the present invention provides a composition for treating a patient in need of anti-angiogenesis therapy comprising a peptide defined above in combination with a pharmaceutically acceptable carrier.

Yet another aspect of the present invention provides a method for treating a patient in need of anti-angiogenesis therapy comprising administering to the patient a therapeutically effective amount of a peptide as defined above.

Still yet another aspect of the present invention provides a composition for the treatment of a disease selected from cancer, arthritis, psoriasis, angiogenesis of the eye associated with infection or surgical intervention, macular degeneration and diabetic retinopathy comprising a peptide as defined above in combination with a pharmaceutically acceptable carrier.

In yet another aspect, the present invention provides a method of isolating a receptor from an endothelial cell comprising binding a peptide as defined above to the receptor to form a peptide receptor complex, isolating the peptide receptor complex, and purifying the receptor.

Detailed Description of the Invention

Definition of Terms

The term "alkyl" as used herein refers to a monovalent group derived from a straight or branched chain saturated hydrocarbon by the removal of a hydrogen atom. Examples of alkyl include, but are not limited to, methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, pentyl, hexyl, and the like. Preferred alkyl groups for the invention are C₁-C₆ alkyl groups having from one to six carbon atoms. Alkyl groups of one to three carbon atoms (C₁-C₃ alkyl) are more preferred for the invention.

The term "nicotiny" as used herein refers to the acyl group derived from nicotinic acid, i.e. pyridine-3-carboxylic acid. The term "2-Me-nicotiny" or "2-methylnicotiny" refers to a nicotiny moiety substituted with a methyl group at the carbon adjacent to the nitrogen atom.

The term "shikimyl" as used herein refers to the acyl residue derived from shikimic acid or [3R-(3 α ,4 α ,5 β)-3,4,5-trihydroxy]-1-cyclohexene-1-carboxylic acid. A "dihydroshikimyl" group denotes the fully saturated analog of shikimic acid.

The term "succinyl" as used herein refers to the acyl residue derived from succinic acid or (1,4-dioxobutyl)-1-carboxylic acid

The term "N-acetylamino" as used herein refers to an amino moiety (-NH₂) substituted on the nitrogen atom with an acetyl (CH₃C(O)-) group.

The term "carbonyl" as used herein refers to the group -C(O)-.

The term "carboxy" or "carboxyl" as used herein refers to the group -C(O)OH.

The term "alkoxy" as used herein refers to an alkyl group as defined above attached to a parent molecular moiety via an ether linkage. Exemplary alkoxy groups include, but are not limited to, methoxy, ethoxy, isopropoxy, and the like.

The term "aromatic ring" as used herein refers to an unsaturated cyclic hydrocarbon associated with a system of π -electron bonds. One to two carbon atoms of the hydrocarbon ring can be substituted with a heteroatom selected from nitrogen, oxygen, or sulfur. Exemplary 5- or 6-membered aromatic rings include, but are not limited to, benzyl, pyridyl, furyl, tetrahydrofuryl, thienyl, and pyrrolyl. An aromatic ring, including rings substituted with a heteroatom, can be optionally substituted on one or more carbon atoms with substituents selected from alkyl, alkoxy, carboxy, and halogen, for example, tolyl, bromobenzyl, *t*-butylbenzyl, nicotiny, 2-methylnicotiny, 2-furoic acid, and the like.

The term "nonaromatic ring" as used herein refers to a saturated or unsaturated cyclic hydrocarbon ring, which can be optionally substituted with one or two heteroatoms

selected from nitrogen, oxygen, or sulfur. Exemplary nonaromatic rings are cyclohexyl, tetrahydropyranyl, pyrrolidinyl, and piperidinyl.

The term "N-protecting group" as used herein refers to an easily removable group which is known in the art to protect an amino group against undesirable reaction during synthetic procedures and to be selectively removable. The use of N-protecting groups is well known in the art for protecting groups against undesirable reactions during a synthetic procedure and many such protecting groups are known, cf, for example, T.H. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, 2nd edition, John Wiley & Sons, New York (1991). Examples of N-protecting groups include, but are not limited to, acyl groups including acetyl, trifluoroacetyl, acylisothiocyanate, aminocaproyl, benzoyl and the like, and acyloxy groups, including *t*-butoxycarbonyl (Boc) and carbobenzyloxy (Cbz), 9-fluorenylmethoxycarbonyl (Fmoc), and the like.

As used herein the terms "Leu," "Sar," "Gln," "Gly," "Val," "Ile," "Thr," "Nva," "Arg," "Asn," "pyroGlu," "Ser," "Ala," "Homoala," "Cha," "Pro," "Phe," "Trp," "1-Nal," "2-Nal," "Azagly" and "Nle" refer to leucine, sarcosine (N-methylglycine), glutamine, glycine, valine, isoleucine, threonine, norvaline, arginine, asparagine, pyroglutamic acid, serine, alanine, homoalanine, cyclohexylalanine, proline, phenylalanine, tryptophan, 1-naphthylalanine, 2-naphthylalanine, azaglycine, and norleucine, respectively, in their L-, D- or DL forms. Unless indicated otherwise by a "D" prefix, *e.g.* D-Ala or D-Ile (also D-Ile), the stereochemistry of the α -carbon of the amino acids and aminoacyl residues in peptides described in this specification and the appended claims is the natural or "L" configuration. The Cahn-Ingold-Prelog "R" and "S" designations are used to specify the stereochemistry of chiral centers in certain of the acyl substituents at the N-terminus of the peptides of this invention. The designation "R,S" is meant to indicate a racemic mixture of the two enantiomeric forms. This nomenclature follows that described in R.S. Cahn, *et al.*, *Angew. Chem. Int. Ed. Engl.*, **5**, 385-415 (1966).

For the most part, the names on naturally occurring and non-naturally occurring aminoacyl residues used herein follow the naming conventions suggested by the IUPAC Commission on the Nomenclature of Organic Chemistry and the IUPAC-IUB Commission on Biochemical Nomenclature as set out in "Nomenclature of α -Amino Acids (Recommendations, 1974)" *Biochemistry*, **14**(2), (1975). To the extent that the names and abbreviations of amino acids and aminoacyl residues employed in this specification and appended claims differ from those suggestions, they will be made clear to the reader. Some abbreviations useful in describing the invention are defined below in the following Table 1.

Table 1

Abbreviation	Definition
Abu	2-aminobutyric acid
6-Ac-Aca	6-N-Ac-caproyl, 6-N-Ac-(CH ₂) ₅ C(O)-, or 6-N-acetyl-aminocaproic acid
Aib	2-aminoisobutyric acid
Ala(3-guanidino)	alanine(3-guanidino)
Ala(3-pyrrolidinylamidino)	alanine[3-pyrrolidinyl(2-N-amidino)]
Ala[4-Pip(N-amidino)]	alanine[4-piperidinyl(N-amidino)]
Allylgly	2-(allyl)glycine
AM	aminomethyl
Aminopyrimidinobutanoyl	2-amino-4-[(2-amino)pyrimidinyl]butanoic acid
Azagly	azaglycine
3-Ac-Bala	3-N-acetyl-beta-alanine
Bala	beta-alanine
Cha	3-(cyclohexyl)alanine
Cha(4-Nisp)	3-(cyclohexyl)alanine(4-N'-isopropyl)
Cit	citrulline
2ClTrt	2-chloro-trityl
Cys(tBu)	cysteine(<i>S-t</i> -butyl)
D-2-Thienylala	D-3-(2-thienyl)alanine
D-3,3-Diphenylala	D-3,3-(diphenyl)alanine
D-3,4-diClPhe	D-3-(3,4-dichlorophenyl)alanine
D-3,4-diFPhe	D-3-(3,4-difluorophenyl)alanine
D-3-Benzothienylala	D-3-(3-benzothienyl)alanine

Table 1 (continued)

D-3-CF ₃ Phe	D-3-(3-trifluoromethylphenyl)alanine
D-3-ClPhe	D-3-(3-chlorophenyl)alanine
D-3-CNPh	D-3-(3-cyanophenyl)alanine
D-3-Pal	D-(3-pyridyl)alanine
D-4,4'-Biphenylala	D-3-(4,4'-biphenyl)alanine
D-4-ClPhe	D-3-(4-chloro-phenyl)alanine
D-Cha	D-3-(cyclohexyl)alanine
D-Chg	D-cyclohexylglycine
Dehydroleu	dehydroleucine
D-Hphe	D-homophenylalanine
D-Ile	D-isoleucine
D-allolIe	D-allo-isoleucine
D-Lys(Nic)	D-lysine(N-epsilon-nicotinyl)
D-Leu	D-leucine
D-pentaFPhe	D-3-(pentafluorophenyl)alanine
D-Val	D-valine
4-Ac-Gaba	4-N-acetyl-gamma-aminobutyric acid or 4-N-acetyl-4-aminobutyric acid
Gaba	gamma-aminobutyric acid or 4-aminobutyric acid
Gly[4-Pip(N-amidino)]	glycine[4-piperidinyI(N-amidino)]
Harg	homoarginine
Hle	homoleucine
Hser	homoserine
Hyp	4-hydroxyproline
Isp	isopropyl
Lys(Ac)	lysine(N-epsilon-acetyl)
Lys(Isp)	lysine(N-epsilon-isopropyl)

Table 1 (continued)

Lys(Nic)	lysine(N-epsilon-nicotinyl)
Met(O)	methionine sulfoxide
Met(O ₂)	methionine sulfone
MeOAc or (MeO)acetyl	methoxyacetyl
1Nal	3-(naphth-1-yl)alanine
2Nal	3-(naphth-2-yl)alanine
N-Ac-Sar	N-acetylsarcosine
Neopentylgly	neopentylglycine
NEtGly	N-ethylglycine
Norarg	norarginine
Octylgly	2-(octyl)glycine
Orn(Ac)	ornithine(N-delta-acetyl)
Orn(2-imidazo)	ornithine [N-delta-(2-imidazolynyl)]
Orn(Isp)	ornithine(N-delta-isopropyl)
Orn(Nic)	ornithine(N-delta-nicotinyl)
O-TBDMS	<i>O</i> - <i>t</i> -butyldimethylsilyl
Pen	penicillamine or β,β -dimethylcysteine
Pen(Acm)	penicillamine(acetamidomethyl)
D-Phe(3,4,5-triF)	D-3-(3,4,5-trifluorophenyl)alanine
D-Phe(3,4-diMeO)	D-3-(3,4-dimethoxyphenyl)alanine
Phe(4-CH ₂ OH)	3-(4-hydroxymethylphenyl)alanine
Phe(4-CONH ₂)	3-(4-carboxyamidophenyl)alanine
Phe(4-guanidino)	3-(4-guanidinophenyl)alanine
D-Phe(4-Me)	D-3-(4-methylphenyl)alanine
D-Phe(4-NH ₂)	D-3-(4-aminophenyl)alanine
Phe(4-NIsp)	3-(4-amino-N-isopropylphenyl)alanine

Table 1 (continued)

Phe(4-CH ₂ NHIsP)	[(4-amino(N-isopropyl)methyl)phenyl]alanine
D-Phe(4-NO ₂)	D-3-(4-nitrophenyl)alanine
Propargylgly	propargylglycine
Pip	pipecolic acid or homoproline
pyBrop	bromo-tris-pyrrolidinophosphoniumhexafluorophosphate
Ser(Bzl)	serine(<i>O</i> -benzyl)
tButylgly	<i>t</i> -butylglycine
Thr(Bzl)	threonine(<i>O</i> -benzyl)
Tic	1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid
Trt	trityl
Tyr(Bzl)	tyrosine(<i>O</i> -benzyl)
Tyr(Et)	tyrosine(<i>O</i> -ethyl)
THF	tetrahydrofuryl or tetrahydrofuran
2-THFcarbonyl	(tetrahydro-2-furyl)carbonyl

When not found in the table above, nomenclature and abbreviations may be further clarified by reference to the Calbiochem-Novabiochem Corp. *1999 Catalog and Peptide Synthesis Handbook* or the Chem-Impex International, Inc. *Tools for Peptide & Solid Phase Synthesis 1998-1999 Catalogue*.

The term "pharmaceutically acceptable salt" as used herein refers to salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, **1977**, 66: 1-19. The salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or separately by reacting the free base function with a suitable organic acid. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphersulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate,

ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydriodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like.

As used herein, the term "pharmaceutically acceptable ester" refers to esters which hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanolic, alkenolic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Examples of particular esters include formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.


The term "pharmaceutically acceptable solvate" represents an aggregate that comprises one or more molecules of the solute, such as a formula (I) compound, with one or more molecules of solvent.

The term "pharmaceutically acceptable prodrugs" as used herein refers to those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

The term "receptor" as used herein refers to a chemical group or molecule on the cell surface or in the cell interior that has an affinity for a specific chemical group,

molecule, or virus. Isolation of receptors relevant to the antiangiogenic activity of the peptide of the invention can provide useful diagnostic tools.


In one embodiment, the present invention relates to compounds of the structure



wherein A_0 , A_1 , A_2 , A_3 , A_7 , A_8 , A_9 , and A_{10} are as defined above. The N-terminus of a nonapeptide represented by A_1 - A_9 can be modified by an amino acyl group represented by A_0 . A_{10} represents a group suitable for modifying the C-terminus of the compound.

10 In the present embodiment, A_4 is an amino acyl residue having a D configuration selected from D-allo-isoleucyl, D-allylglycyl, D-3-(3-cyanophenyl)alanyl, D-cystyl, D-isoleucyl, D-leucyl, D-penicillaminy, D-phenylalanyl, D-3-(3,4,5-trifluorophenyl)alanyl, and D-3-(4-aminophenyl)alanyl; A_5 is an amino acyl residue selected from octylglycyl, glycyl, penicillaminy, seryl, threonyl, and tyrosyl; and A_6 is an amino acyl residue selected from glutaminy, leucyl, norvalyl, and seryl.

15 In another embodiment of the invention, the compounds have the structure (I) as defined above wherein A_1 is sarcosyl, A_2 is glycyl, A_3 is valyl, A_7 is isoleucyl, A_8 is arginyl, and A_9 is prolyl. Compounds of the present embodiment can be represented by the structure



20 wherein A_0 is hydrogen or an acyl group modifying the N-terminus. Suitable groups for A_0 can be represented by the formula $R\text{-(CH}_2\text{)}_n\text{-C(O)-}$; wherein n is an integer from 0 to 8 and R is selected from hydroxyl; methyl; N-acetylamino; methoxyl; carboxyl; cyclohexyl optionally containing a one or two double bonds and optionally substituted with one to three hydroxyl groups; and a 5- or 6-membered ring aromatic or nonaromatic ring
25 optionally containing one or two heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the ring is optionally substituted with a moiety selected from alkyl, alkoxy, and halogen; or $R^1\text{-CH}_2\text{CH}_2\text{-(OCH}_2\text{CH}_2\text{O)}_p\text{-CH}_2\text{-C(O)-}$; wherein R^1 is selected from hydrogen, alkyl, and N-acetylamino, and p is an integer from 1 to 8.


A_4 is an amino acyl residue of L or D configuration selected from allo-isoleucyl,
30 dehydroleucyl, glycyl, isoleucyl, prolyl, D-alanyl, D-3-(naphth-1-yl)alanyl, D-3-(naphth-

2-yl)alanyl, D-(3-pyridyl)-alanyl, D-2-aminobutyryl, D-allo-isoleucyl, D-allo-threonyl, D-allylglycyl, D-asparaginy, D-aspartyl, D-benzothienyl, D-3-(4,4'-biphenyl)alanyl, D-chlorophenylalanyl, D-3-(3-trifluoromethylphenyl)alanyl, D-3-(3-cyanophenyl)alanyl, D-3-(3,4-difluorophenyl)alanyl, D-citrullyl, D-cyclohexylalanyl, D-cyclohexylglycyl, D-cystyl, D-cystyl(*S*-*t*-butyl), D-glutaminy, D-glutamyl, D-histidyl, D-homoisoleucyl, D-homophenylalanyl, D-homoseryl, D-isoleucyl, D-leucyl, D-lysyl(N-epsilon-nicotinyl), D-lysyl, D-methionyl, D-neopentylglycyl, D-norleucyl, D-norvalyl, D-ornithyl, D-penicillaminy, D-penicillaminy(acetamidomethyl), D-penicillaminy(*S*-benzyl), D-phenylalanyl, D-3-(4-aminophenyl)alanyl, D-3-(4-methylphenyl)alanyl, D-3-(4-nitrophenyl)alanyl, D-3-(3,4-dimethoxyphenyl)alanyl, D-3-(3,4,5-trifluorophenyl)alanyl, D-prolyl, D-seryl, D-seryl(*O*-benzyl), D-*t*-butylglycyl, D-thienylalanyl, D-threonyl, D-threonyl(*O*-benzyl), D-tryptyl, D-tyrosyl(*O*-benzyl), D-tyrosyl(*O*-ethyl), D-tyrosyl, and D-valyl.

A₅ is an amino acyl residue of L or D configuration selected from alanyl, 3-pyridyl)-alanyl, 3-(naphth-1-yl)alanyl, 3-(naphth-2-yl)alanyl, allo-threonyl, allylglycyl, glutaminy, glycyl, histidyl, homoseryl, isoleucyl, lysyl(N-epsilon-acetyl), methionyl, norvalyl, octylglycyl, ornithyl, 3-(4-hydroxymethylphenyl)alanyl, prolyl, seryl, threonyl, tryptyl, tyrosyl, D-allo-threonyl, D-homoseryl, D-seryl, D-threonyl, penicillaminy, and cystyl.

A₆ is an amino acyl residue of L or D configuration selected from alanyl, 3-(naphth-1-yl)alanyl, 3-(naphth-2-yl)alanyl, (3-pyridyl)alanyl, 2-aminobutyryl, allylglycyl, arginyl, asparaginy, aspartyl, citrullyl, cyclohexylalanyl, glutaminy, glutamyl, glycyl, histidyl, homoalanyl, homoleucyl, homoseryl, isoleucyl, leucyl, lysyl(N-epsilon-acetyl), lysyl(N-epsilon-isopropyl), methionyl(sulfone), methionyl(sulfoxide), methionyl, norleucyl, norvalyl, octylglycyl, phenylalanyl, 3-(4-carboxamidephenyl)alanyl, propargylglycyl, seryl, threonyl, tryptyl, tyrosyl, valyl, D-3-(naphth-1-yl)alanyl, D-3-(naphth-2-yl)alanyl, D-glutaminy, D-homoseryl, D-leucyl, D-norvalyl, D-seryl, penicillaminy, and cystyl.

A₁₀ is a hydroxyl group or an amino acid amide selected from azaglycylamide, D-alanylamide, D-alanylethylamide, glycylamide, glycylethylamide, sarcosylamide, serylamide, D-serylamide, or A₁₀ is a group represented by the formula



or a group represented by the formula $-\text{NH}-\text{R}^4$, wherein s is an integer selected from 0 to 8; R^2 is selected from hydrogen, alkyl, and a 5- to 6-membered cycloalkyl ring; R^3 is selected from hydrogen, hydroxy, alkyl, phenyl, alkoxy, and a 5- to 6-membered ring optionally containing from one to two heteroatoms selected from oxygen, nitrogen, and sulfur, provided that s is not zero when R^3 is hydroxy or alkoxy; and R^4 is selected from hydrogen, hydroxy, and a 5- to 6-membered cycloalkyl ring.

Preferred compounds of the invention have the structure (II) as defined above, wherein A_4 is an amino acyl residue having a D configuration selected from D-alanyl, D-3-(naphth-1-yl)alanyl, D-3-(naphth-2-yl)alanyl, D-(3-pyridyl)-alanyl, D-2-aminobutyryl, D-allo-isoleucyl, D-allo-threonyl, D-allylglycyl, D-asparaginy, D-aspartyl, D-chlorophenylalanyl, D-3-(3-trifluoromethylphenyl)alanyl, D-3-(3-cyanophenyl)alanyl, D-3-(3,4-difluorophenyl)alanyl, D-cyclohexylalanyl, D-cyclohexylglycyl, D-cystyl, D-glutaminy, D-glutamyl, D-histidyl, D-homoisoleucyl, D-homophenylalanyl, D-homoseryl, D-isoleucyl, D-leucyl, D-lysyl(N-epsilon-nicotinyl), D-methionyl, D-neopentylglycyl, D-norleucyl, D-norvalyl, D-penicillaminy, D-penicillaminy(acetamidomethyl), D-penicillaminy(*S*-benzyl), D-phenylalanyl, D-3-(4-aminophenyl)alanyl, D-3-(4-methylphenyl)alanyl, D-3-(4-nitrophenyl)alanyl, D-3-(3,4-dimethoxyphenyl)alanyl, D-3-(3,4,5-trifluorophenyl)alanyl, D-prolyl, D-seryl, D-seryl(*O*-benzyl), D-*t*-butylglycyl, D-thienylalanyl, D-threonyl, D-threonyl(*O*-benzyl), D-tyrosyl(*O*-ethyl), D-tyrosyl, D-valyl, and D-cystyl.

Other preferred compounds of the present invention have the structure of formula (II), wherein A_5 is selected from glycyl, octylglycyl, penicillaminy, seryl, threonyl, and tyrosyl.

Additional preferred compounds of the present invention have the structure represented by formula (II), wherein A_6 is selected from glutaminy, leucyl, norvalyl, and seryl.

The more preferred amino acid residues for substituting the position represented by A_4 are D configuration amino acids selected from D-allo-isoleucyl, D-allylglycyl, D-3-(3-cyanophenyl)alanyl, D-cystyl, D-isoleucyl, D-leucyl, D-penicillaminy, D-phenylalanyl, D-3-(3,4,5-trifluorophenyl)alanyl, and D-3-(4-aminophenyl)alanyl.

Preferred A₀ groups for modifying the N-terminus of the compounds in the scope of the invention are selected from acetyl, butyryl, caproyl, (4-N-acetylamino)butyryl, N-acetyl-beta-alanyl, (6-N-acetylamino)caproyl, chloronicotinyl, cyclohexylacetyl, furoyl, gamma-aminobutyryl, 2-methoxyacetyl, methylnicotinyl, nicotinyl, (8-N-acetylamino)-
5 3,6-dioxo-octanoyl, phenylacetyl, propionyl, shikimyl, succinyl, and tetrahydrofuroyl.

The preferred A₁₀ groups for modifying the C-terminus of the invention are selected from D-alanylamine, azaglycylamine, serylamine, ethylamine, hydroxylamine, isopropylamine, propylamine, 2-(cyclohexyl)ethylamine, 2-(1-pyrrolidine)ethylamine, 1-(cyclohexyl)ethylamine, 2-(methoxy)ethylamine, 2-(hydroxy)ethylamine, 2-(2-pyridine)ethylamine, (2-pyridine)methylamine, 2-(3-pyridine)ethylamine, 2-(2-(1-methyl)pyrrolidine)ethylamine, 2-(N-morpholine)ethylamine, and cyclopropylmethylamine.
10

Compounds contemplated as falling within the scope of the present invention include, but are not limited to:

15 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
pyroGlu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
20 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂-(1-pyrrolidine),
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethylpiperidine,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHmethylcyclopropyl,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH(ethyl-1-(R)-cyclohexyl),
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂,
25 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂cyclohexyl,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
30 N-Ac-Sar-Gly-Val-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-Gly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Val-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Met-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Nle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Sar-Gly-Val-D-Tyr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-4,4'-Biphenylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cha-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Chg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-4-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Hphe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-Dehydroleu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-3-CF₃Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-pentaFPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-3,4-diClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-3-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-2-Thienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-3-CNPh-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-DNva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Ile-Thr-Cha-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gly-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Abu-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-Ile-Thr-Allylgly-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Octylgly-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Met-Ile-Arg-ProNHCH₂CH₃,
 N-Cyclohexylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Nicotinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Propionyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-(MeO)acetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Butyryl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-[CH₃CONH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃
 N[6-N-acetyl-(CH₂)₅C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 N-Hexanoyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-[4-N-Acetylaminobutyryl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 H-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Asn-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Pro-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Gly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Ala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-NEtGly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-AbuNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Phe-NHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Tic-NHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Hyp-NHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Aib-NHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-Ala-NHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pip-NHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Tyr(Et)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys(tBu)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Tyr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ser(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-1Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-tButylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Sar-Gly-Val-D-Orn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Thr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-2Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(4-Me)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe-(3,4-diMeO)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-(4-NO₂)Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen(Acm)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Abu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-Phe(4-NH₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ala-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Met-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Phe-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Tyr-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Nva-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Asp-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Gly-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Lys(Ac)-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Leu-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-2Nal-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-1Nal-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Allylgly-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Cit-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ala-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-Leu-Pro-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Trp-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Leu-Nva-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Lys(Ac)-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-2Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Sar-Gly-Val-D-Leu-1Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Octylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Gln-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Met-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Leu-Allylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ile-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-D-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ile-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nle-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-Ile-Thr-Cit-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O₂)-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Arg-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Tyr-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Glu-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Ile-Thr-Lys(Ac)-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Propargylgly-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Bala-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Phenylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-Azagly-NH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Sar-NHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SerNH₂,
 N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Leu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Phe-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Glu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Pro-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Asn-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Asp-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Asn-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Gln-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Ser-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Glu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Gaba-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Bala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Gln-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Gly-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Glu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Asp-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Ile-Thr-Asp-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Asn-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O)-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Asn-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cit-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-Hcit-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Hle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Neopentylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- N-Ac-Sar-Gly-Val-D-Ile-Thr-Phe(4-CONH₂)-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-His-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Isp)-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Nic)-ProNHCH₂CH₃,
5 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(Nic)-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(Isp)-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-NIsp)-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cha(4-NIsp)-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Harg-ProNHCH₂CH₃,
10 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Norarg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cit-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Phe(4-CH₂OH)-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-guanidino)-ProNHCH₂CH₃,
15 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Aminopyrimidinylbutanoyl-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-CH₂NH(Isp))-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Gly[4-Pip(N-amidino)]-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala[4-Pip(N-amidino)]-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-guanidino)-ProNHCH₂CH₃,
20 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-pyrrolidinylamidino)-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(2-imidazo)-ProNHCH₂CH₃,
N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
25 N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
30 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SarNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-SarNH₂,
 5 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-SarNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-SarNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Ile-Thr-Orn(Ac)-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂,
 N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 20 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 25 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 30 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 5 N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 10 N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 15 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂,
 20 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 25 N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-AzaglyNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHethyl-(1-pyrrolidine),
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNH(ethyl-1-cyclohexyl),
 30 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHethyl-(1-pyrrolidine),
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl),
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl),

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₂OCH₃,
 5 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 10 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Allygly-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 15 N-Ac-Sar-Gly-Val-D-Ile-Ser-Allygly-Ile-Arg-Pro-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Allygly-Ile-Arg-Pro-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SarNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-NHOH,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Hser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Gln-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Nva-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Ile-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Phe-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Leu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Ser-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Thr-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ala-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,

- N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Ala-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Ala-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Val-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Val-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Leu-Ser-Val-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-D-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
 15 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-D-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-D-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 25 N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 30 N-Ac-Sar-Gly-Val-D-Leu-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 5 N-Ac-Sar-Gly-Val-D-alloIle-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Ile-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-Ile-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Arg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-3-Pal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Glu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Asp-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-His-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloThr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-D-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ser-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-Thr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloThr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ser-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

N-Ac-Sar-Gly-Val-D-Thr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloThr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloThr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Thr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 5 N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 10 N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 15 N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 20 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl-1-(S)-cyclohexyl,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-Pen-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-Pen-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Pen-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 5 N-Succinyl-Sar-Gly-Val-D-Pen-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Cys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Cys-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-Cys-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cys-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 20 N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Cys-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Pen-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Pen-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Pen-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 N-Succinyl-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Leu-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,

- N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Pen-Ser-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Sar-Gly-Val-D-Ile-Pen-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂CH₃,
 10 N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-Pro-D-AlaNH₂,
 15 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Pen-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Leu-Gly-Pen-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Pen-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 25 N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Ala-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Ala-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Ala-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Ala-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Ala-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-Pro-D-AlaNH₂,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 10 N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-(3-Ac-Bala)-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Ala-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Ala-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Ala-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 N-(3-Ac-Bala)-Sar-Ala-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-Pro-OH,
 25 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-OH,
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-Pro-OH,
 30 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-Pro-OH
 N-Ac-Sar-Gly-Asp-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Ala-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Cha-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Met-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Hser-Ile-Arg-ProNHCH₂CH₃,
 5 N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-alloIle-His-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-Butyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Butyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Amyl,
 10 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-hexyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3,3-dimethyl)butyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-ethoxy)ethyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-isopropoxy)ethyl,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3-methoxy)propyl,
 15 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(cyclopentyl)methyl, and
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-cyclohexyl.

Preferred compounds for the practice of the invention are:

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂-(1-pyrrolidine),
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH(ethyl-1-(R)-cyclohexyl),
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 N-Ac-Sar-Gly-Val-D-Val-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Nle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Cha-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-3,4-diClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 N-Ac-Sar-Gly-Val-D-3-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-2-Thienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-3-CNPh-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- N-Ac-Sar-Gly-Val-D-Ile-Thr-Cha-Ile-Arg-ProNHCH₂CH₃,
N[2-THFcarbonyl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N[6-N-acetyl-(CH₂)₅C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Hexanoyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
5 N-[4-N-Acetylaminobutyl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Pro-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-NEtGly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
10 N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Lys(Ac)-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Leu-Arg-ProNHCH₂CH₃,
15 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-1Nal-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Allylgly-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Ala-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Trp-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
20 N-Ac-Sar-Gly-Val-D-Leu-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-2Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-1Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Octylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
25 N-Ac-Sar-Gly-Val-D-Leu-Allylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Leu-D-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Tyr-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Glu-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Propargylgly-Ile-Arg-ProNHCH₂CH₃,
30 N-Ac-Sar-Gly-Val-D-allol-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Bala-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Phenylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂,
N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SerNH₂,

N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 5 N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 10 N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃, and
 N-Ac-Sar-Gly-Val-D-Phe(4-NH₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃.

It is well known in the art that modifications and changes can be made in the
 20 structure of a polypeptide without substantially altering the biological function of that
 peptide. For example, certain amino acids can be substituted for other amino acids in a
 given polypeptide without any appreciable loss of function. In making such changes,
 substitutions of like amino acid residues can be made on the basis of relative similarity of
 side-chain substituents, for example, their size, charge, hydrophobicity, hydrophilicity,
 25 and the like.

In describing the invention, certain abbreviations are used for the sake of
 convenience throughout the specification, including the examples, to refer to reagents and
 compounds useful for preparing the compounds of the invention. When so used, the
 following abbreviations are meant to refer to the following: DMF for dimethylformamide;
 30 DMA for dimethylacetamide; DIEA for diisopropylethylamine; HATU for O-(7-aza-
 benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate; NMP for N-
 methylpyrrolidone; and TFA for trifluoroacetic acid.

Determination of Biological Activity

Pellet Preparation

Ten microliters of a mixture containing a final concentration of 1, 5, or 10 mM of the peptides of invention, 100 ng of bFGF (Collaborative Biomedical Products, Bedford, MA), and 6% Hydron (Sigma, St. Louis, MO) were pipetted into the tip of a sterile Teflon rod. After drying for 1-2 hours, the pellets were stored at 4 °C.

Pellet Implantation

A small (about 2 mm) radial incision at 1 mm from the center of the cornea was performed in anesthetized Sprague Dawley rats. With a curved iris spatula, an intrastromal pocket was made to a distance of 1 mm from the limbus-the circular blood vessels that surround the cornea. A single pellet was implanted. Antibiotic ointment (neosporin) was applied post surgery to the operated eye to prevent infection and to decrease inflammation.

Data Analysis

At day seven post-implantation, neovascularization was measured through a slitlamp biomicroscopy (Nikon NS-1), connected to an image analysis system (Leica Qwin). The response was calculated by colorimetrically detecting the area of new blood vessels, and calculating the new vessel surface area in μm^2 . The compounds of the invention inhibit rat cornea neovascularization as shown in Table 2.

Table 2
Effect of the Compounds of the Invention on Rat Cornea Neovascularization

<u>Peptide</u>	<u>Number of Corneas/Dose</u>	<u>% Inhibition</u>
Example 1	6/10 μM	92.6
Example 1	5/5 μM	74.8
Example 1	4/6 μM	71.5
untreated	5/-	-

The compounds of the invention, including but not limited to those specified in the examples, possess anti-angiogenic activity. As angiogenesis inhibitors, such compounds are useful in the treatment of both primary and metastatic solid tumors, including carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach, pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney, bladder and urothelium), female genital tract, (including cervix, uterus, and ovaries as well as choriocarcinoma and gestational trophoblastic disease), male genital tract (including

prostate, seminal vesicles, testes and and germ cell tumors), endocrine glands (including the thyroid, adrenal, and pituitary glands), and skin, as well as hemangiomas, melanomas, sarcomas (including those arising from bone and soft tissues as well as Kaposi's sarcoma) and tumors of the brain, nerves, eyes, and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas, and meningiomas). Such compounds may also be useful in treating solid tumors arising from hematopoietic malignancies such as leukemias (i.e. chloromas, plasmacytomas and the plaques and tumors of mycosis fungoides and cutaneous T-cell lymphoma/leukemia) as well as in the treatment of lymphomas (both Hodgkin's and non-Hodgkin's lymphomas).

In addition, these compounds may be useful in the prevention of metastases from the tumors described above either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

Further uses include the treatment and prophylaxis of autoimmune diseases such as rheumatoid, immune and degenerative arthritis; various ocular diseases such as diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, retrolental fibroplasia, neovascular glaucoma, rubeosis, retinal neovascularization due to macular degeneration, hypoxia, angiogenesis in the eye associated with infection or surgical intervention, and other abnormal neovascularization conditions of the eye; skin diseases such as psoriasis; blood vessel diseases such as hemangiomas, and capillary proliferation within atherosclerotic plaques; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; and wound granulation. Other uses include the treatment of diseases characterized by excessive or abnormal stimulation of endothelial cells, including but not limited to intestinal adhesions, Crohn's disease, atherosclerosis, scleroderma, and hypertrophic scars, i.e. keloids.

Another use is as a birth control agent, by inhibiting ovulation and establishment of the placenta. The compounds of the invention are also useful in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochelia minimalis quintosa*) and ulcers (*Helicobacter pylori*). The compounds of the invention are also useful to reduce bleeding by administration prior to surgery, especially for the treatment of resectable tumors.

The compounds of the invention may be used in combination with other compositions and procedures for the treatment of diseases. For example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with a peptide of the present invention and then a peptide of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize and inhibit the growth of any residual primary tumor. Additionally, the compounds of the

invention may be combined with pharmaceutically acceptable excipients, and optionally sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions.

A sustained-release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid-base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix desirably is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid) polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

When used in the above or other treatments, a therapeutically effective amount of one of the compounds of the present invention may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt form. By a "therapeutically effective amount" of the compound of the invention is meant a sufficient amount of the compound to treat an angiogenic disease, (for example, to limit tumor growth or to slow or block tumor metastasis) at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate,

camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethansulfonate (isothionate), lactate, maleate, methanesulfonate, nicotine, 2-naphthalenesulfonate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, phosphate, glutamate, bicarbonate, *p*-toluenesulfonate and undecanoate. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as acetic acid, maleic acid, succinic acid and citric acid. Other salts include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic basis. Preferred salts of the compounds of the invention include phosphate, tris and acetate.

Alternatively, a compound of the present invention may be administered as pharmaceutical compositions containing the compound of interest in combination with one or more pharmaceutically acceptable excipients. A pharmaceutically acceptable carrier or excipient refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The compositions may be administered parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), rectally, or buccally. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions for parenteral injection comprise pharmaceutically-acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also

be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters), poly(anhydrides), and (poly)glycols, such as PEG. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Topical administration includes administration to the skin or mucosa, including surfaces of the lung and eye. Compositions for topical administration, including those for inhalation, may be prepared as a dry powder which may be pressurized or non-pressurized. In non-pressurized powder compositions, the active ingredient in finely divided form may be used in admixture with a larger-sized pharmaceutically-acceptable inert carrier comprising particles having a size, for example, of up to 100 micrometers in diameter. Suitable inert carriers include sugars such as lactose. Desirably, at least 95% by weight of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 micrometers.

Alternatively, the composition may be pressurized and contain a compressed gas, such as nitrogen or a liquified gas propellant. The liquified propellant medium and indeed the total composition is preferably such that the active ingredient does not dissolve therein to any substantial extent. The pressurized composition may also contain a surface active agent, such as a liquid or solid non-ionic surface active agent or may be a solid anionic surface active agent. It is preferred to use the solid anionic surface active agent in the form of a sodium salt.

A further form of topical administration is to the eye. A compound of the invention is delivered in a pharmaceutically acceptable ophthalmic vehicle, such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye, as for example the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically-acceptable ophthalmic vehicle may, for example, be an ointment, vegetable oil or an

encapsulating material. Alternatively, the compounds of the invention may be injected directly into the vitreous and aqueous humour.

Compositions for rectal or vaginal administration are preferably suppositories which may be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 *et seq.*

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they may also be used in combination with one or more agents which are conventionally administered to patients for treating angiogenic diseases. For example, the compounds of the invention are effective over the short term to make tumors more sensitive to traditional cytotoxic therapies such as chemicals and radiation. The compounds of the invention also enhance the effectiveness of existing cytotoxic adjuvant anti-cancer therapies. The compounds of the invention may also be combined with other antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In particular, when used in the treatment of solid tumors, compounds of the invention may be administered with IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopril, angioinhibins, TNP-470, pentosan polysulfate, platelet factor 4, LM-609, SU-5416, CM-101, Tecogalan, plasminogen-K-5, vasostatin, vitaxin, vasculostatin, squalamine, marimastat or other MMP inhibitors, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, cisplatin,

taxol, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, and the like as well as with radiation.

Total daily dose of the compositions of the invention to be administered to a human or other mammal host in single or divided doses may be in amounts, for example, from 0.0001 to 300 mg/kg body weight daily and more usually 1 to 300 mg/kg body weight.

It will be understood that agents which can be combined with the compound of the present invention for the inhibition, treatment or prophylaxis of angiogenic diseases are not limited to those listed above, but include in principle any agents useful for the treatment or prophylaxis of angiogenic diseases.

The peptides of the invention may be used for the development of affinity columns for isolation of receptors relevant to the antiangiogenic activity of the peptide of the invention, e.g. TSP-1 receptor, in, for example, cultured endothelial cells. As is known in the art, isolation and purification of the receptor may be followed by amino acid sequencing to identify and isolate polynucleotides which encode the receptor. Recombinant expression of this receptor would allow greater amounts of receptor to be produced, e.g. to produce a sufficient quantity for use in high throughput screening assays to identify other angiogenesis inhibitors.

The peptides of the present invention may be chemically coupled to isotopes, enzymes, carrier proteins, cytotoxic agents, fluorescent molecules, chemiluminescent, bioluminescent and other compounds for a variety of applications. For example, a peptide may be labeled to facilitate testing of its ability to bind antisera or to detect cell types which possess a relevant receptor. The coupling technique is generally chosen on the basis of the functional groups available on the amino acids of the peptide including, but not limited to amino, sulfhydryl, carboxyl, amide, phenol, and imidazole. Various reagents used to effect such couplings include among others, glutaraldehyde, diazodized benzidine, carbodiimide, and p-benzoquinone.

The efficiency of the coupling reaction is determined using different techniques appropriate for the specific reaction. For example, radiolabeling of the peptide with ^{125}I may be accomplished using chloramine T and NaI^{125} of high specific activity. The reaction is terminated with sodium metabisulfite and the mixture is desalted on disposable columns. The labeled peptide is eluted from the column and fractions are collected. Aliquots are removed from each fraction and radioactivity measured in a gamma counter. In this manner, a labeled peptide may be obtained which is free from unreacted NaI^{125} .

The peptides of the present invention can also be used as antigens to generate polyclonal or monoclonal antibodies. Such antibodies can be used in diagnostic methods

and kits to detect or quantify the peptide of the invention, or peptides related thereto, in a body fluid or tissue. Results from these tests could be used to diagnose or determine the prognostic relevance of such peptides.

The use of the peptides of the present invention to generate monoclonal antibodies in animals such as the mouse, rabbit or sheep, follows techniques well known in the art. If desired, the antibodies can then be used to make anti-idiotypic antibodies which in turn can be humanized as is known in the art to prevent immunological responses. The humanized antibodies can be used to inhibit angiogenesis or to make kits to detect the receptor as described herein.

For the production of polyclonal antisera in rabbits, sheep, goats or other animals the peptides of the invention are coupled, for example through lysine residues, to purified bovine serum albumin using glutaraldehyde. The efficiency of this reaction may be determined by measuring the incorporation of radiolabeled peptide. Unreacted glutaraldehyde and peptide may be separated by dialysis and the conjugate stored for subsequent use.

Serum samples from generation of polyclonal antisera or media samples from production of monoclonal antisera may be analyzed for determination of antibody titer and in particular, for the determination of high titer antisera. Subsequently, the highest titer antisera may be tested to establish the following: a) optimal antiserum dilution for highest specific binding of the antigen and lowest non-specific binding, b) ability to bind increasing amounts of peptide in a standard displacement curve, c) potential cross-reactivity with immunologically-related peptides and proteins (including plasminogen, TSP-1, and TSP-1 of related species), and d) ability to detect the peptide of the invention in extracts of plasma, urine, tissues, and in cell culture media.

Titer may be established through several means known in the art, such as by dot blot and density analysis, and also by precipitation of radiolabeled peptide-antibody complexes using protein A, secondary antisera, cold ethanol or charcoal-dextran followed by activity measurement with a gamma counter. If desired, the highest titer antisera may be purified on affinity columns. For example, the peptides of the invention may be coupled to a commercially available resin and used to form an affinity column. Antiserum samples may then be passed through the column so that antibodies to the peptides of the invention bind (via the peptide) to the column. These bound antibodies are subsequently eluted, collected and evaluated for determination of titer and specificity.

Kits for measurement of the compounds of the invention are also contemplated as part of the present invention. Antisera that possess the highest titer and specificity and can detect the peptides of the invention in extracts of plasma, urine, tissues, and in cell culture

media may be used to establish assay kits for rapid, reliable, sensitive, and specific measurement and localization of peptides of the invention. These assay kits may employ (but are not limited to) the following techniques: competitive and non-competitive assays, radioimmunoassay (RIA), bioluminescence and chemiluminescence assays, fluorometric assays, sandwich assays, immunoradiometric assays, dot blots, enzyme linked assays including ELISA, microtiter plates, antibody coated strips or dipsticks for rapid monitoring of urine or blood, and immunocytochemistry. For each kit the range, sensitivity, precision, reliability, specificity and reproducibility of the assay are established by means well known to those skilled in the art.

The above described assay kit would provide instructions, antiserum, one or more peptides of the invention, and possibly radiolabeled peptides of the invention and/or reagents for precipitation of bound peptide/antibody complexes. Such a kit would be useful for the measurement of the peptide of the invention in biological fluids and tissue extracts of animals and humans with and without tumors, as is well known in the art.

Another kit may be used to visualize or localize the peptide of the invention in tissues and cells. Immunohistochemistry techniques and kits, for example, which employ such techniques are well known to those of ordinary skill in the art. Such a kit provides antisera to the peptide of the invention, and possibly blocking serum and secondary antiserum linked to a fluorescent molecule such as fluorescein isothiocyanate, or to some other reagent used to visualize the primary antiserum. Using this methodology, biopsied tumors may be examined for sites of peptide production or for sites of the peptide receptor. Alternatively, a kit may supply radiolabeled nucleic acids for use in *in situ* hybridization to probe for messenger RNA which encodes the compound of the invention.

Synthesis of the Peptides

The polypeptides of the present invention may be synthesized by any techniques that are known to those skilled in the art. For solid phase peptide synthesis, a summary of the many techniques may be found in J.M. Stewart and J.D. Young, Solid Phase Peptide Synthesis, W.H. Freeman Co. (San Francisco), 1963 and J. Meienhofer, Hormonal Proteins and Peptides, vol. 2, p. 46, Academic Press (New York), 1973. For classical solution synthesis see G. Schroder and K. Lupke, The Peptides, vol. 1, Academic Press (New York), 1965.

Reagents, resins, amino acids, and amino acid derivatives are commercially available and can be purchased from Chem-Impex International, Inc. (Wood Dale, IL, U.S.A.) or Calbiochem-Novabiochem Corp. (San Diego, CA, U.S.A.) unless otherwise noted herein.

In general, these methods comprise the sequential addition of one or more amino acids or suitably protected amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid is protected by a suitable protecting group. The protected or derivatized amino acid can then be either attached to an inert solid support or utilized in solution by adding the next amino acid in the sequence having the complimentary (amino or carboxyl) group suitably protected, under conditions suitable for forming the amide linkage. The protecting group is then removed from this newly added amino acid residue and the next amino acid (suitably protected) is then added, and so forth. After all the desired amino acids have been linked in the proper sequence, any remaining protecting groups (and any solid support) are removed sequentially or concurrently, to afford the final polypeptide. By simple modification of this general procedure, it is possible to add more than one amino acid at a time to a growing chain, for example, by coupling (under conditions which do not racemize chiral centers) a protected tripeptide with a properly protected dipeptide to form, after deprotection, a pentapeptide.

A particularly preferred method of preparing compounds of the present invention involves solid phase peptide synthesis.

In this particularly preferred method the alpha-amino function is protected by an acid or base sensitive group. Such protecting groups should have the properties of being stable to the conditions of peptide linkage formation, while being readily removable without destruction of the growing peptide chain or racemization of any of the chiral centers contained therein. Suitable protecting groups are 9-fluorenylmethyloxycarbonyl (Fmoc), *t*-butyloxycarbonyl (Boc), benzyloxycarbonyl (Cbz), biphenylisopropyl-oxycarbonyl, *t*-amyloxycarbonyl, isobornyloxycarbonyl, (α,α)-dimethyl-3,5-dimethoxybenzyloxycarbonyl, *o*-nitrophenylsulfonyl, 2-cyano-*t*-butyloxycarbonyl, and the like. The 9-fluorenylmethyloxycarbonyl (Fmoc) protecting group is preferred.

Particularly preferred side chain protecting groups are, for side chain amino groups as in lysine and arginine: 2,2,5,7,8-pentamethylchroman-6-sulfonyl (pmc), nitro, *p*-toluenesulfonyl, 4-methoxybenzenesulfonyl, Cbz, Boc, and adamantyloxycarbonyl; for tyrosine: benzyl, *o*-bromobenzyloxycarbonyl, 2,6-dichlorobenzyl, isopropyl, *t*-butyl (*t*-Bu), cyclohexyl, cyclopentyl and acetyl (Ac); for serine: *t*-butyl, benzyl and tetrahydropyranyl; for histidine: trityl, benzyl, Cbz, *p*-toluenesulfonyl and 2,4-dinitrophenyl; for tryptophan: formyl and Boc.

In the solid phase peptide synthesis method, the C-terminal amino acid is attached to a suitable solid support or resin. Suitable solid supports useful for the above synthesis are those materials which are inert to the reagents and reaction conditions of the stepwise condensation-deprotection reactions, as well as being insoluble in the media used. The

preferred solid support for synthesis of C-terminal carboxy peptides is 4-hydroxymethyl-phenoxymethyl-copoly(styrene-1% divinylbenzene). The preferred solid support for C-terminal amide peptides is 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)phenoxy-acetamidoethyl resin available from Applied Biosystems.

The C-terminal amino acid is coupled to the resin by means of N,N'-dicyclohexylcarbodiimide (DCC), N,N'-diisopropylcarbodiimide (DIC) or O-benzotriazol-1-yl-N,N,N',N'-tetramethyluroniumhexafluorophosphate (HBTU), with or without 4-dimethylaminopyridine (DMAP), 1-hydroxybenzotriazole (HOBT), benzotriazol-1-yloxy-tris(dimethylamino)phosphoniumhexafluorophosphate (BOP) or bis(2-oxo-3-oxazolidinyl)phosphine chloride (BOPCl), mediated coupling for from about 1 to about 24 hours at a temperature of between 10 ° and 50 °C in a solvent such as dichloromethane or DMF. When the solid support is 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamidoethyl resin, the Fmoc group is cleaved with a secondary amine, preferably piperidine, prior to coupling with the C-terminal amino acid as described above. The preferred method for coupling to the deprotected 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)phenoxyacetamidoethyl resin is O-benzotriazol-1-yl-N,N,N',N'-tetramethyluroniumhexafluorophosphate (HBTU, 1 equiv.) and 1-hydroxybenzotriazole (HOBT, 1 equiv.) in DMF.

The coupling of successive protected amino acids can be carried out in an automatic polypeptide synthesizer as is well known in the art. In a preferred embodiment, the α -amino function in the amino acids of the growing peptide chain are protected with Fmoc. The removal of the Fmoc protecting group from the N-terminal side of the growing peptide is accomplished by treatment with a secondary amine, preferably piperidine. Each protected amino acid is then introduced in about 3-fold molar excess and the coupling is preferably carried out in DMF. The coupling agent is normally O-benzotriazol-1-yl-N,N,N',N'-tetramethyluroniumhexafluorophosphate (HBTU, 1 equiv.) and 1-hydroxybenzotriazole (HOBT, 1 equiv.).

At the end of the solid phase synthesis, the polypeptide is removed from the resin and deprotected, either in succession or in a single operation. Removal of the polypeptide and deprotection can be accomplished in a single operation by treating the resin-bound polypeptide with a cleavage reagent, for example thianisole, water, ethanedithiol and trifluoroacetic acid.

In cases wherein the C-terminus of the polypeptide is an alkylamide, the resin is cleaved by aminolysis with an alkylamine. Alternatively, the peptide may be removed by transesterification, e.g. with methanol, followed by aminolysis or by direct transamidation. The protected peptide may be purified at this point or taken to the next step directly. The

removal of the side chain protecting groups is accomplished using the cleavage cocktail described above.

The fully deprotected peptide is purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivatized polystyrene-divinylbenzene (for example, AMBERLITE® XAD); silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g. on SEPHADEX® G-25, LH-20 or countercurrent distribution; high performance liquid chromatography (HPLC), especially reverse-phase HPLC on octyl- or octadecylsilyl-silica bonded phase column packing.

The following examples will serve to further illustrate the preparation of the novel compounds of the invention.

Preparation of the Cleavage Reagent

The cleavage reagent (2 mL) is prepared by mixing, in the following order, thioanisole (100 µL), water (50 µL), ethanedithiol (50 µL) and trifluoroacetic acid (1.8 mL). The freshly-prepared mixture is cooled to -5 °C to -10 °C and used as described below.

Cleavage and Deprotection Procedure

A mixture of resin-bound polypeptide and cleavage reagent is stirred at 0 °C for 10-15 minutes and then at ambient temperature for a further 1.75 hours. The amount of time is increased by 0.5 hours for each additional arginine up to a total of three hours. The amount of cleavage reagent used is determined using the following formula:

<u>weight of resin (mg)</u>	<u>amount of cleavage reagent (µL)</u>
0-10	100
10-25	200
25-50	400
50-100	700
100-200	1200

The resin is then filtered off and rinsed with neat trifluoroacetic acid. The filtrate is then added in 0.5 mL portions to a centrifuge tube containing about 8 mL of cold diethyl ether. The suspension is then centrifuged and the supernatant is decanted off. The pellet is re-suspended in about 8 mL of ether, another 0.5 mL of the filtrate is added, and the

process is repeated until all of the peptide is precipitated. The precipitated filtrate is then washed with ether, dried and lyophilized.

- If the peptide does not precipitate upon addition to ether, the mixture is shaken with aqueous 30% acetic acid. The organic phase is then extracted twice with aqueous 30% acetic acid and the combined aqueous extracts are lyophilized.

Example 1

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- In the peptide synthesis column position of a Perkin Elmer/Applied Biosynthesis SYNERGY® peptide synthesizer is placed an Pro(2-CITrt) peptide synthesis column (25 μM amino acid; Nova Biochem). Amino acids are added sequentially according to the following synthetic cycle:

- (1) Solvating the resin using DMF for about 5 minutes;
- (2) Washing with DMF for about 5 minutes;
- (3) Activating the incoming Fmoc protected amino acid (75 μM) using a 0.2 M solution of HBTU (75 μM) and HOBT (75 μM) in DMSO-NMP (N-methylpyrrolidone);
- (4) Coupling using a solution in DMF of the activated Fmoc protected amino acid prepared in step 3 above for about 30 minutes;
- (5) Washing with DMF for 5 minutes; and
- (6) For peptides capped with acetyl at the N-terminus, substituting acetic acid (87 μM) for an Fmoc protected amino acid and using 87 μM each of HBTU and HOBT.
- (7) For peptides capped with ethylamide at the C-terminus, adding DMF to the resin followed by ByProp (1.1 equivalents) and ethylamine (20 equivalents) in THF.

The amino acids were coupled to the resin in the following order using the conditions indicated.

# Amino Acid	Coupling
1. Fmoc-Arg(Pmc)	30 minutes
2. Fmoc-Ile	30 minutes
3. Fmoc-Nva	30 minutes
4. Fmoc-Thr(<i>t</i> -Bu)	30 minutes
5. Fmoc-D-Ile	30 minutes
6. Fmoc-Val	30 minutes
7. Fmoc-Gly	30 minutes

8. Fmoc-Sar	30 minutes
-------------	------------

Upon completion of the synthesis, the resin was washed with THF for about 5 minutes to remove DMF and shrink the resin. The resin was then gas dried with argon for about 10 minutes and nitrogen for a further 10 minutes to provide the resin-bound peptide (85 mg). Cleavage and deprotection are accomplished using the procedure described above (40 mg of dry resin-bound peptide, 700 μ L of cleavage reagent, cleavage time 2.5 hours) to give the crude peptide (14 mg). Purification by HPLC using a 7 μ m Symmetry Prep C18 column (7.8x300 mm) with solvent mixtures varying in a gradient from 5% to 100% acetonitrile-water over a period of 50 minutes followed by lyophilization provided the desired peptide.

The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 26.5 min (10% to 40% acetonitrile in water containing 0.01% of TFA, over 30 min period); MS (ESI) m/e 994 (M+H)⁺.

Example 2

pyroGlu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

# Amino Acid	Coupling
1. Fmoc-Arg(Pmc)	30 minutes
2. Fmoc-Ile	30 minutes
3. Fmoc-Nva	30 minutes
4. Fmoc-Thr(<i>t</i> -Bu)	30 minutes
5. Fmoc-D-Ile	30 minutes
6. Fmoc-Val	30 minutes
7. Fmoc-Gly	30 minutes
8. pyroGlu(Boc)	30 minutes

The desired peptide was prepared using the conditions described for Example 1. The amino acids were coupled to the resin in the following order using the conditions indicated.

The pure fractions were lyophilized to yield pyroGlu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 23.5 min (gradient of 10% to 40% acetonitrile in water containing 0.01% of TFA, over 30 min period); MS (ESI) m/e 994 (M+H)⁺.

Example 3

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₃

The procedure described in Example 1 was used but substituting methylamine (2.0 M solution in THF) for ethylamine. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₃ as the trifluoroacetate salt: R_t = 3.224 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 930 (M+H)⁺; Amino Acid Anal.: 1.09 Sar; 1.03 Gly; 0.98 Val; 0.98 Ile; 0.54 Thr; 1.72 Nva; 1.01 Arg; 1.08 Pro.

Example 4

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 1 was used but substituting isopropylamine for ethylamine. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHisopropyl as the trifluoroacetate salt: R_t = 3.648 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1008 (M+H)⁺; Amino Acid Anal.: 1.10 Sar; 0.99 Gly; 0.96 Val; 1.88 Ile; 0.56 Thr; 1.67 Nva; 0.96 Arg; 1.09 Pro.

Example 5

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl-(1-pyrrolidine)

Resin Preparation

4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin (0.5 g, 0.54 mmol/g substitution) was placed in a solid phase synthesis reaction vessel containing (9:1) DMA/acetic acid (4 mL). The mixture was shaken for 5 min. The resin was drained and this process was repeated three times. To the swollen resin were added 10-15 grains of activated 4A molecular sieves and (9:1) DMA/acetic acid (4mL) and 10 molar equivalents of 1-(2-aminoethyl)pyrrolidine. The slurry was shaken for 1h at rt and to it was added 10 molar equivalents of sodium triacetoxyborohydride. The slurry was shaken for 2 h at rt. The resin was drained and washed three times with DMA, three times with methanol, three times with dichloromethane, three times with diethyl ether and dried in vacuo at rt overnight. The dry resin was swollen in DMA (4 mL) and shaken for 5 min. This process was repeated twice.

Coupling of Fmoc-Pro

To the swollen resin in the reaction vessel were added sequentially the following chemicals: DMA (4 mL), one equivalent of DIEA, a DMA solution containing 3.0 equivalents of Fmoc-Pro, 3.0 equivalents of HATU, and 3.0 equivalents of DIEA. The slurry was shaken overnight. The resin was drained and washed three times with DMA, three times with methanol, three times with dichloromethane, three times with diethyl ether and dried in vacuo at rt overnight. A small portion of the resin was used to determine the Fmoc-Pro loading. The rest of the resin was shaken with DMA (4 mL) three times for 5 min and then for 1 h at rt with a solution of (8:1:1) DMA/pyridine/acetic anhydride (5 mL). The resin was drained and washed three times with DMA, three times with methanol, three times with dichloromethane, and three times with diethyl ether. The

resin was dried in vacuo at rt overnight and then used in the subsequent solid phase peptide synthesis.

Synthesis of above peptide

In the synthesis of the above peptide the amino acids, the coupling conditions and the synthetic protocol used were the identical to as those described in Example 1. Upon completion of the synthesis the peptide and the protecting groups were cleaved at rt using (95:5) TFA/anisole (3 mL) for 3h. The resin was filtered and washed three times with methanol. The combined filtrates were concentrated in vacuo and to the residue was added diethylether. The solid precipitate was filtered. The crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl(1-pyrrolidine) as the bis-trifluoroacetate salt: $R_t = 4.40$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1063 ($\text{M}+\text{H}^+$); Amino Acid Anal.: 0.95 Sar; 1.0 Gly; 0.86 Val; 1.63 Ile; 0.56 Thr; 1.38 Nva; 0.88 Arg; 1.07 Pro.

Example 6

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl(1-piperidine)

The procedure described in Example 5 was used but substituting 1-(2-aminoethyl)piperidine for 1-(2-aminoethyl)pyrrolidine in the reductive alkylation step. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl-(1-piperidine) as the bis-trifluoroacetate salt: $R_t = 4.437$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1077

(M+H)⁺; Amino Acid Anal.: 1.11 Sar; 1.04 Gly; 0.99 Val; 1.77 Ile; 0.61 Thr; 1.61 Nva; 0.97 Arg; 1.10 Pro.

Example 7

5 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHmethylcyclopropyl

The procedure described in Example 1 was used but substituting (aminoethyl)cyclopropane for 1-(2-aminoethylpyrrolidine). After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHmethylcyclopropyl as the trifluoroacetate salt: R_t = 3.815 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1020 (M+H)⁺; Amino Acid Anal.: 1.01 Sar; 0.96 Gly; 0.96 Val; 1.66 Ile; 0.53 Thr; 1.65 Nva; 1.08 Arg; 1.09 Pro.

15

Example 8

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl

The procedure described in Example 5 was used but substituting (R)-1-cycloylethylamine for 1-(2-aminoethylpyrrolidine). After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate salt: R_t = 5.196 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1076 (M+H)⁺; Amino Acid Anal.: 1.19 Sar; 0.99 Gly; 0.62 Val; 1.47 Ile; 0.48 Thr; 1.57 Nva; 1.01 Arg; 0.83 Pro.

25

Example 9

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH(2-hydroxyethyl)


The procedure described in Example 5 was used but substituting O-TBDMS-ethanolamine for 1-(2-aminoethylpyrrolidine). After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH(2-hydroxyethyl) as the trifluoroacetate salt: $R_t = 4.04$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1010 ($\text{M}+\text{H}$)⁺; Amino Acid Anal.: 1.04 Sar; 1.01 Gly; 0.98 Val; 1.59 Ile; 0.44 Thr; 1.45 Nva; 0.99 Arg; 1.06 Pro.

Example 10

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂


The procedure described in Example 1 was used but substituting Fmoc-Pro-Sieber amide resin for H-Pro-2-CITrt resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL), the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂ as the trifluoroacetate salt: $R_t = 4.063$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 966 ($\text{M}+\text{H}$)⁺; Amino Acid Anal.: 0.87 Sar; 0.98 Gly; 0.94 Val; 1.73 Ile; 0.47 Thr; 1.35 Nva; 1.02 Arg; 1.05 Pro.

Example 11



The procedure described in Example 5 was used but substituting 2-methoxyethylamine for 1-(2-aminoethylpyrrolidine). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂-OCH₃ as the trifluoroacetate salt: R_t = 3.40 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1024 (M+H)⁺; Amino Acid Anal.: 1.02 Sar; 1.06 Gly; 0.97 Val; 1.54 Ile; 0.47 Thr; 1.81 Nva; 0.97 Arg; 1.25 Pro.

Example 12



The procedure described in Example 5 was used but substituting cyclohexylethylamine for 1-(2-aminoethylpyrrolidine). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂-cyclohexyl as the trifluoroacetate salt: R_t = 4.97 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1076 (M+H)⁺; Amino Acid Anal.: 0.87 Sar; 1.00 Gly; 0.88 Val; 1.34 Ile; 0.44 Thr; 1.61 Nva; 1.07 Arg; 1.05 Pro.

Example 13

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂CH₃

The procedure described in Example 1 was used but substituting propylamine for ethylamine. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂CH₃ as the trifluoroacetate salt: R_t = 3.68 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1008 (M+H)⁺; Amino Acid Anal.: 0.94 Sar; 1.09 Gly; 0.96 Val; 1.58 Ile; 0.51 Thr; 1.78 Nva; 0.96 Arg; 1.23 Pro.

Example 14

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 22.5 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺; Amino Acid Anal.: 0.95 Sar; 0.96 Gly; 0.97 Val; 0.99 Ile; 0.54 Thr; 1.66 Nva; 1.14 Arg; 1.08 Pro.

Example 15

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.54 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺; Amino Acid Anal.: 1.00 Sar; 0.93 Gly; 0.96 Val; 1.02 Leu; 0.58 Thr; 1.50 Nva; 0.99 Ile; 1.14 Arg; 1.08 Pro.

Example 16

N-Ac-Sar-Gly-Val-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ile for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.28 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺; Amino Acid Anal.: 0.95 Sar; 0.94 Gly; 0.89 Val; 1.70 Ile; 0.52 Thr; 1.67 Nva; 0.99 Ile; 1.27 Arg; 1.06 Pro.

Example 17

N-Ac-Sar-Gly-Val-Gly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gly for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-Gly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 2.47$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 938 (M+H)⁺; Amino Acid Anal.: 1.10 Sar; 1.94 Gly; 1.03 Val; 0.98 Ile; 0.54 Thr; 1.61 Nva; 1.28 Arg; 1.05 Pro.

Example 18

N-Ac-Sar-Gly-Val-D-Val-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Val for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Val-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.13$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 980 (M+H)⁺; Amino Acid Anal.: 1.07 Sar; 1.0 Gly; 2.01 Val; 0.99 Ile; 0.62 Thr; 1.54 Nva; 1.49 Arg; 1.11 Pro.

Example 19

N-Ac-Sar-Gly-Val-allolle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-allolle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-allolle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.174 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺; Amino Acid Anal.: 1.02 Sar; 0.99 Gly; 0.95 Val; 1.29 Ile; 0.45 Thr; 1.52 Nva; 1.54 Arg; 1.07 Pro.

Example 20

N-Ac-Sar-Gly-Val-D-Ala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Ala for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.826 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 952 (M)⁺ and 908 (M-44)⁺.

Example 21

N-Ac-Sar-Gly-Val-D-Lys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Lys(Boc) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the

protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Lys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt:

- 5 $R_t = 3.544$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1009 (M)⁺ and 965 (M-44)⁺.

Example 22

N-Ac-Sar-Gly-Val-D-Met-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- 10 The procedure described in Example 1 was used but substituting Fmoc-D-Met for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Met-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.141$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1012 (M)⁺.
- 15

Example 23

N-Ac-Sar-Gly-Val-D-Nle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- 20 The procedure described in Example 1 was used but substituting Fmoc-D-Nle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Nle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.383$ min
- 25

(gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period);
MS (APCI) m/e 994 (M)⁺.

Example 24

N-Ac-Sar-Gly-Val-D-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Phe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.476 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1028 (M)⁺.

Example 25

N-Ac-Sar-Gly-Val-D-Trp-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Trp(Boc) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Trp-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.430 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1024 (M)⁺.

Example 26

N-Ac-Sar-Gly-Val-D-Tyr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Tyr(2-CITrt) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Tyr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.964 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1045 (M)⁺.

Example 27

N-Ac-Sar-Gly-Val-D-4,4'-Biphenylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-4,4'-Biphenylala for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-4,4'-Biphenylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.005 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1104 (M)⁺.

Example 28

N-Ac-Sar-Gly-Val-D-Cha-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Cha for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Cha-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.005 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period);

5 MS (APCI) m/e 1034 (M)⁺.

Example 29

N-Ac-Sar-Gly-Val-D-Chg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Chg for


10 Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Chg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.377 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period);
15 MS (APCI) m/e 977 (M)⁺.

Example 30

N-Ac-Sar-Gly-Val-D-4-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

20 The procedure described in Example 1 was used but substituting Fmoc-D-4-ClPhe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield
25 N-Ac-Sar-Gly-Val-D-4-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.674 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1018 (M)⁺.


Example 31



The procedure described in Example 1 was used but substituting Fmoc-D-Hphe for


- 5 Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Hphe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.597 min
10 (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1042 (M)⁺ and 998 (M-44)⁺.

Example 32



- 15 The procedure described in Example 1 was used but substituting Fmoc-Dehydroleu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield
20 N-Ac-Sar-Gly-Val-Dehydroleu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.1707 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 992 (M)⁺ and 949 (M-44)⁺.

Example 33



- 25 The procedure described in Example 1 was used but substituting Fmoc-D-3-CF₃Phe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the

protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-3-CF₃Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.825 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1097 (M)⁺ and 1053 (M-44)⁺.

Example 34

N-Ac-Sar-Gly-Val-D-pentaFPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-pentaFPhe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-pentaFPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.810 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1118 (M)⁺ and 1075 (M-44)⁺.

Example 35

N-Ac-Sar-Gly-Val-D-3,4-diClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-3,4-diClPhe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield

N-Ac-Sar-Gly-Val-D-3,4-diCIPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 4.911 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1100 (M+3)⁺.

5

Example 36

N-Ac-Sar-Gly-Val-D-3-CIPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- 10 The procedure described in Example 1 was used but substituting Fmoc-D-3-CIPhe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield
- 15 N-Ac-Sar-Gly-Val-D-3-CIPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 4.689 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1062 (M)⁺.

Example 37

- 20 N-Ac-Sar-Gly-Val-D-2-Thienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-D-2-Thienylala for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to
- 25 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-2-Thienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the

trifluoroacetate salt: $R_t = 4.388$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1034 (M)⁺.

Example 38

5 N-Ac-Sar-Gly-Val-D-3-CNPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-3-CN-Phe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield
10 N-Ac-Sar-Gly-Val-D-3-CNPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.361$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1009 (M)⁺.

15 Example 39

N-Ac-Sar-Gly-Val-D-3,3-Diphenylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-3,3'-Diphenylala for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by
20 C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-3,3'-Diphenylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.778$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1104 (M)⁺.

25

Example 40

N-Ac-Sar-Gly-Val-D-3-Benzothienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-3-Benzothienylala for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-3-Benzothienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.797 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1084 (M)⁺.

Example 41

N-Ac-Sar-Gly-Val-D-3,4-diF-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-3,4-diF-Phe for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-3,4-diF-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.608 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1064 (M)⁺.

Example 42

N-Ac-Sar-Gly-Val-D-Ile-Thr-DNva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-DNva for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-

water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-DNva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.75 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺; Amino Acid Anal.: 1.08 Sar; 0.96 Gly; 0.95 Val; 1.74 Ile; 5 0.50 Thr; 1.69 Nva; 1.26 Arg; 1.09 Pro.

Example 43

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gln(Trt) 10 for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.047 min 15 (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1023 (M+H)⁺; Amino Acid Anal.: 1.15 Sar; 0.96 Gly; 0.63 Val; 1.7 Ile; 0.46 Thr; 0.65 Glu; 1.45 Arg; 1.04 Pro.

Example 44

N-Ac-Sar-Gly-Val-D-Ile-Thr-Cha-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Cha for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile- 25 water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Cha-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.503 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period);

MS (ESI) m/e 1048 (M+H)⁺; Amino Acid Anal.: 1.18 Sar; 0.94 Gly; 0.59 Val; 1.65 Ile; 0.45 Thr; 0.37 Cha; 1.45 Arg; 1.06 Pro.

Example 45

5 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gly-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gly for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-
10 water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gly-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.11 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 952 (M+H)⁺.


Example 46

15 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ala for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column
20 chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.16 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 966 (M+H)⁺.


25

Example 47




The procedure described in Example 1 was used but substituting Fmoc-Val for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.36 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺.

Example 48



The procedure described in Example 1 was used but substituting Fmoc-Abu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Abu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.23 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 980 (M+H)⁺.

Example 49



The procedure described in Example 1 was used but substituting Fmoc-Allylgly for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting

groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Allylgly-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.40 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 992 (M+H)⁺.

Example 50

N-Ac-Sar-Gly-Val-D-Ile-Thr-Octylgly-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Octylgly for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Octylgly-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.30 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1064 (M+H)⁺.

Example 51

N-Ac-Sar-Gly-Val-D-Ile-Thr-Met-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Met for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Met-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.48 min

(gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period);
MS (ESI) m/e 1027 (M+H)⁺.

Example 52

5 N-Cyclohexylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting cyclohexylacetic acid for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Cyclohexylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.11 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1076 (M+H)⁺; Amino Acid Anal.: 1.15 Sar; 0.97 Gly; 0.95 Val; 1.79 Ile; 0.54 Thr; 1.66 Nva; 1.28 Arg; 1.08 Pro.

15

Example 53

N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting 2-Me-nicotinic acid for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.11 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1071 (M+H)⁺; Amino Acid Anal.: 1.19 Sar; 1.01 Gly; 0.99 Val; 1.79 Ile; 0.57 Thr; 1.70 Nva; 1.59 Arg; 1.17 Pro.

25

Example 54

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but acylating the peptide resin (after the Fmoc-Sar coupling and deprotection) with a (1:1) succinic anhydride/pyridine mixture (2 mL) overnight. After washing the resin and cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.72 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1052 (M+H)⁺; Amino Acid Anal.: 1.16 Sar; 1.05 Gly; 0.95 Val; 1.85 Ile; 0.57 Thr; 1.70 Nva; 1.59 Arg; 1.17 Pro.

Example 55

N-Nicotinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting nicotinic acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Nicotinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.6 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1057 (M+H)⁺; Amino Acid Anal.: 1.03 Sar; 0.89 Gly; 0.81 Val; 1.48 Ile; 0.40 Thr; 1.46 Nva; 1.07 Arg; 1.04 Pro.

Example 56

N-Propionyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting propionic acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Propionyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.7 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M+H)⁺; Amino Acid Anal.: 0.93 Sar; 0.97 Gly; 0.88 Val; 1.60 Ile; 0.44 Thr; 1.58 Nva; 1.17 Arg; 1.10 Pro.

Example 57

N-MeOacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting methoxyacetic acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-MeOacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.45 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M+H)⁺; Amino Acid Anal.: 1.12 Sar; 1.06 Gly; 0.94 Val; 1.62 Ile; 0.48 Thr; 1.91 Nva; 1.40 Arg; 1.27 Pro.

Example 58

N-Shikimyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting shikimic acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Shikimyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.0 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1108 (M+H)⁺; Amino Acid Anal.: 1.22 Sar; 1.06 Gly; 0.94 Val; 1.80 Ile; 0.55 Thr; 1.70 Nva; 1.28 Arg; 1.26 Pro.

Example 59

N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting 2-furoic acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.0 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1046 (M+H)⁺; Amino Acid Anal.: 1.02 Sar; 1.00 Gly; 0.99 Val; 1.66 Ile; 0.45 Thr; 1.75 Nva; 1.45 Arg; 1.21 Pro.

Example 60

N-Butyryl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃


The procedure described in Example 1 was used but substituting butyric acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Butyryl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.03 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1022 (M+H)⁺; Amino Acid Anal.: 1.13 Sar; 0.99 Gly; 1.01 Val; 1.93 Ile; 0.67 Thr; 1.61 Nva; 1.45 Arg; 1.08 Pro.

Example 61

N-(Tetrahydro-2-furoyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃


The procedure described in Example 1 was used but substituting tetrahydro-2-furoic acid for acetic acid at the last coupling. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-(tetrahydro-2-furoyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.91 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1050 (M+H)⁺; Amino Acid Anal.: 1.12 Sar; 0.97 Gly; 0.88 Val; 1.41 Ile; 0.42 Thr; 1.60 Nva; 1.43 Arg; 1.03 Pro.

Example 62



The procedure described in Example 1 was used but coupling with Fmoc-8-amino-3,6-dioxo-octanoic acid after the Fmoc-Sar coupling, after removal of the terminal Fmoc the peptide resin was coupled with acetic acid as described above. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.32 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1139 (M+H)⁺; Amino Acid Anal.: 1.04 Sar; 1.01 Gly; 0.91 Val; 1.67 Ile; 0.53 Thr; 1.77 Nva; 1.39 Arg; 1.02 Pro.

Example 63



The procedure described in Example 1 was used but coupling with Fmoc-6-amino-hexanoic acid after the Fmoc-Sar coupling, after removal of the terminal Fmoc the peptide resin was coupled with acetic acid as described above. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-[6-N'-Acetyl-(CH₂)₅C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.60 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI)

m/e 1107 (M+H)⁺; Amino Acid Anal.: 1.13 Sar; 0.96 Gly; 0.89 Val; 1.42 Ile; 0.43 Thr; 1.68 Nva; 1.44 Arg; 1.04 Pro.

Example 64

5 N-Hexanoyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting hexanoic acid for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-
10 water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Hexanoyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 4.95 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1050 (M+H)⁺; Amino Acid Anal.: 1.07 Sar; 0.93 Gly; 1.02 Val; 1.95 Ile; 0.56 Thr; 1.31 Nva; 1.52 Arg; 1.05 Pro.

15

Example 65

N-[4-N'-Acetyl-buteryl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but coupling with Fmoc-4-amino-butyrlic acid after the Fmoc-Sar coupling, after removal of the terminal Fmoc the peptide
20 resin was coupled with acetic acid as described above. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-[4-N'-Acetyl-buteryl]-Sar-Gly-Val-D-Ile-Thr-Nva-
25 Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 4.09 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1079

(M+H)⁺; Amino Acid Anal.: 1.03 Gaba; 1.07 Sar; 0.93 Gly; 1.00 Val; 1.90 Ile; 0.54 Thr; 1.30 Nva; 1.54 Arg; 1.06 Pro.

Example 66

H-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃


The procedure described in Example 1 was used but omitting the acetic acid coupling at the end. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield H-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the bistrifluoroacetate salt: R_t = 3.65 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 952 (M+H)⁺; Amino Acid Anal.: 1.00 Sar; 1.00 Gly; 0.99 Val; 1.67 Ile; 0.50 Thr; 1.76 Nva; 1.47 Arg; 1.22 Pro.

Example 67

N-Ac-Sar-Gly-Asn-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃


The procedure described in Example 1 was used but substituting Fmoc-Asn(Trt) for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Asn-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the bistrifluoroacetate salt: R_t = 2.45 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1009 (M+H)⁺; Amino Acid Anal.: 1.05 Sar; 0.98 Gly; 0.96 Asp; 1.7 Ile; 0.48 Thr; 1.54 Nva; 1.32 Arg; 1.07 Pro.

Example 68



The procedure described in Example 1 was used but substituting Fmoc-8-amino-3,6-dioxo-octanoic acid for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.12 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1068 (M+H)⁺; Amino Acid Anal.: 0.93 Gly; 1.02 Val; 1.97 Ile; 0.57 Thr; 1.31 Nva; 1.54 Arg; 1.05 Pro.

Example 69



The procedure described in Example 1 was used but substituting Fmoc-Pro for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Pro-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.30 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1020 (M+H)⁺; Amino Acid Anal.: 0.92 Gly; 0.99 Val; 1.80 Ile; 0.50 Thr; 1.32 Nva; 1.53 Arg; 2.09 Pro.

Example 70

N-Ac-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gly for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.08 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 980 (M+H)⁺; Amino Acid Anal.: 1.89 Gly; 1.02 Val; 1.91 Ile; 0.52 Thr; 1.35 Nva; 1.57 Arg; 1.09 Pro.

Example 71

N-Ac-Ala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ala for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Ala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.00 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺; Amino Acid Anal.: 1.01 Ala; 0.93 Gly; 1.01 Val; 1.92 Ile; 0.56 Thr; 1.30 Nva; 1.51 Arg; 1.05 Pro.

Example 72

N-Ac-NEtGly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-NEtGly for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-NEtGly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.24 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M+H)⁺; Amino Acid Anal.: 0.95 Gly; 1.04 Val; 1.99 Ile; 0.59 Thr; 1.34 Nva; 1.50 Arg; 1.01 Pro.

Example 73

N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Leu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.348 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M+H)⁺; Amino Acid Anal.: 0.88 Sar; 0.99 Gly; 0.95 Val; 1.03 Ile; 0.55 Thr; 1.12 Leu; 1.53 Arg; 1.07 Pro.

Example 74

N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield

N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt:

R_t = 3.963 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 982 (M+H)⁺; Amino Acid Anal.: 0.91 Sar; 0.97 Gly; 1.00 Val; 1.03 Ile; 0.56 Thr; 0.23 Ser; 1.52 Arg; 1.08 Pro.

Example 75

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 10 was used but substituting Fmoc-D-Ala-Siebert amide resin for Fmoc-Pro-Siebert amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt: R_t = 4.117 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1037 (M+H)⁺; Amino Acid Anal.: 0.85 Sar; 0.94 Gly; 0.92 Val; 1.83 Ile; 0.54 Thr; 1.18 Nva; 1.01 Arg; 1.04 Pro; 1.01 Ala.

Example 76

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-ProNHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-D-Pro-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-Pro-NHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.20 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M+H)⁺.

Example 77

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-AbuNHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-Abu-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-AbuNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.35 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 982 (M+H)⁺.

Example 78

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-PheNHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-Phe-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the

crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Phe-NHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.73$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1044 (M+H)⁺.

Example 79

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Tic-NHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-Tic-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Tic-NHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.68$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1056 (M+H)⁺.

Example 80

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Hyp-NHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-Hyp-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Hyp-NHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.95$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1010 (M+H)⁺.

Example 81

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Aib-NHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-Aib-

- 5 Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Aib-
- 10 NHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.25 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 982 (M+H)⁺.

Example 82

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-Ala-NHCH₂CH₃

- 15 The procedure described in Example 10 was used but substituting Fmoc-D-Ala-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure
- 20 fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-Ala-NHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.95 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 968 (M+H)⁺.

Example 83

- 25 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pip-NHCH₂CH₃

The procedure described in Example 10 was used but substituting Fmoc-Pip-Sieber ethylamide resin for Fmoc-Pro-Sieber amide resin. After cleavage of the peptide from the

resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pip-NHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.30 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M+H)⁺.

Example 84

N-Ac-Sar-Gly-Val-D-Tyr(Et)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Tyr(Et) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Tyr(Et)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 6.01 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1072 (M)⁺.

Example 85

N-Ac-Sar-Gly-Val-D-Cys(tBu)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Cys(tBu) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Cys(tBu)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.96 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1040 (M)⁺.

Example 86

N-Ac-Sar-Gly-Val-D-Cys(Acm)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Cys(Acm) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Cys(Acm)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.12 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1044 (M)⁺.

Example 87

N-Ac-Sar-Gly-Val-D-Tyr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Tyr(Bzl) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Tyr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 6.74 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1135 (M+H)⁺.

Example 88

N-Ac-Sar-Gly-Val-D-Ser(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Ser(Bzl) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ser(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.95 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1058 (M)⁺.

Example 89

N-Ac-Sar-Gly-Val-D-1Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-1Nal for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-1Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 6.30 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1081 (M+3)⁺.

Example 90

N-Ac-Sar-Gly-Val-D-tButylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-tButylgly for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-tButylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.46 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 994 (M)⁺.

Example 91

N-Ac-Sar-Gly-Val-D-Orn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Orn(Boc) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Orn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 1.69 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 995 (M)⁺.

Example 92

N-Ac-Sar-Gly-Val-D-Thr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Thr(Bzl) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Thr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 6.10 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1072 (M)⁺.

Example 93

N-Ac-Sar-Gly-Val-D-2Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-2Nal for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-2Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 6.33 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (APCI) m/e 1078 (M)⁺.

Example 94

N-Ac-Sar-Gly-Val-D-Phe(4-Me)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Phe(4-Me) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Phe(4-Me)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate

salt: $R_t = 3.654$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1042 (M)⁺.

Example 95

N-Ac-Sar-Gly-Val-D-Phe(3,4-diMeO)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Phe(3,4-diMeO) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Phe(3,4-diMeO)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.006$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1088 (M)⁺.

Example 96

N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Phe(3,4,5-triF) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.848$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1082 (M)⁺.

Example 97

N-Ac-Sar-Gly-Val-D-Phe(4-NO₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Phe(4-NO₂) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-

Ac-Sar-Gly-Val-D-Phe(4-NO₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.483 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1073 (M)⁺.

5

Example 98

N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Pen(Trt) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 2.928 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1012 (M)⁺.

15

Example 99

N-Ac-Sar-Gly-Val-D-Pen(Acm)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Pen(Acm) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Pen(Acm)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 2.415 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1083 (M)⁺.

25

Example 100

N-Ac-Sar-Gly-Val-D-Pen(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Pen(Bzl) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50%

30

acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Pen(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.124 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1102 (M)⁺.

5

Example 101

N-Ac-Sar-Gly-Val-D-Abu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Abu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Abu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.533 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 966 (M)⁺.

Example 102

N-Ac-Sar-Gly-Val-D-Phe(4-NH₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-Phe(4-Boc-NH₂) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Phe(4-NH₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.545 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1043 (M)⁺.

Example 103

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ala-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Ala for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ala-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.675 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period);
5 MS (ESI) m/e 952 (M)⁺.

Example 104

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Gln-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Gln(Trt)
10 for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Gln-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.46 min
15 (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1009 (M)⁺.

Example 105

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Met-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Met for
20 Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Met-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.219 min
25 (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1012 (M)⁺.

Example 106

30 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Phe-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Phe for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting

groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Phe-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.579 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1028 (M)⁺.

Example 107

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Pro-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Pro for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Pro-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.704 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 978 (M)⁺.

Example 108

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ser-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Ser(tBu) for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ser-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.510 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 968 (M)⁺.

Example 109

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Trp-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Trp(Boc) for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Trp-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.625$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1067 (M)⁺.

Example 110

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Tyr-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Tyr(tBu) for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Tyr-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.017$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1044 (M)⁺.

Example 111

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Nva-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Nva for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Nva-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.139$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 980 (M)⁺.

Example 112

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Asp-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-

- 5 Asp(OtBu)-OH for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Asp-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t =
10 2.082 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 996 (M)⁺.

Example 113

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Gly-Arg-ProNHCH₂CH₃

- 15 The procedure described in Example 15 was used but substituting Fmoc-Gly for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-
20 Val-D-Leu-Thr-Nva-Gly-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.623 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 938 (M)⁺.

Example 114

- 25 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Lys(Ac)-Arg-ProNHCH₂CH₃

- The procedure described in Example 15 was used but substituting Fmoc-Lys(Ac) for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-
30 water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Lys(Ac)-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.599

min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1051 (M)⁺.

Example 115

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Leu-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Leu for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Leu-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.403 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M)⁺.

Example 116

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-2Nal-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-2Nal for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-2Nal-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.198 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1078 (M)⁺.

Example 117

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-1Nal-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-1Nal for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-

Val-D-Leu-Thr-Nva-1Nal-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.217 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1078 (M)⁺.

5

Example 118

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Allylgly-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Allylgly for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Allylgly-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.993 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 978 (M)⁺.

15

Example 119

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Cit-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Cit for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Cit-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.408 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1038 (M)⁺.

25

Example 120

N-Ac-Sar-Gly-Val-D-Leu-Ala-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Ala for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50%

30

acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ala-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.481 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 964 (M)⁺.

5

Example 121

N-Ac-Sar-Gly-Val-D-Leu-Pro-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Pro for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Pro-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.621 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 990 (M)⁺.

Example 122

N-Ac-Sar-Gly-Val-D-Leu-Trp-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Trp(Boc) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Trp-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 4.378 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1079 (M)⁺.

Example 123

N-Ac-Sar-Gly-Val-D-Leu-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Tyr(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18

column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.606 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1056 (M)⁺.

Example 124

N-Ac-Sar-Gly-Val-D-Leu-Nva-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Nva for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Nva-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.870 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 992 (M)⁺.

Example 125

N-Ac-Sar-Gly-Val-D-Leu-Gly-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Gly for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Gly-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.397 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 950 (M)⁺.

Example 126

N-Ac-Sar-Gly-Val-D-Leu-Lys(Ac)-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Lys(Ac) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the

protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Lys(Ac)-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.365$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1063 (M)⁺.

Example 127

N-Ac-Sar-Gly-Val-D-Leu-2Nal-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-2Nal for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-2Nal-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 4.992$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1090 (M)⁺.

Example 128

N-Ac-Sar-Gly-Val-D-Leu-1Nal-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-1Nal for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-1Nal-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 5.032$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1090 (M)⁺.

Example 129

N-Ac-Sar-Gly-Val-D-Leu-Octylgly-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Octylgly for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Octylgly-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 5.90 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1062 (M)⁺.

Example 130

N-Ac-Sar-Gly-Val-D-Leu-Gln-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Gln(Trt) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Gln-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.323 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1021 (M)⁺.

Example 131

N-Ac-Sar-Gly-Val-D-Leu-Met-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Met for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Met-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.901 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M)⁺.

Example 132

N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 15 was used but substituting Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.414$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 980 (M)⁺.

Example 133

N-Ac-Sar-Gly-Val-D-Leu-Allylgly-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 15 was used but substituting Fmoc-Allylgly for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Allylgly-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.801$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 990 (M)⁺.

Example 134

N-Ac-Sar-Gly-Val-D-Leu-Ile-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 15 was used but substituting Fmoc-Ile for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ile-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t =$

4.028 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1006 (M)⁺.

Example 135

N-Ac-Sar-Gly-Val-D-Leu-D-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-D-Thr(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-D-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.437 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M)⁺.

Example 136

N-Ac-Sar-Gly-Val-D-Ile-Thr-Ile-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ile for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Ile-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.54 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M)⁺; Amino Acid Anal.: 1.07 Sar; 0.94 Gly; 0.91 Val; 3.02 Ile; 0.47 Thr; 1.24 Arg; 1.04 Pro.

Example 137

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nle-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Nle for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-

water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nle-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.80 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1006 (M)⁺.

5

Example 138

N-Ac-Sar-Gly-Val-D-Ile-Thr-Cit-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Cit for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Cit-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.83 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1052 (M)⁺; Acid Anal.: 1.05 Sar; 1.00 Gly; 1.00 Val; 2.13 Ile; 0.65 Thr; 1.11 Cit; 1.49 Arg; 1.10 Pro.

15

Example 139

N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O₂)-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Met(O₂) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O₂)-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.701 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1058 (M)⁺; Acid Anal.: 1.36 Sar; 0.94 Gly; 0.62 Val; 2.06 Ile; 0.13 Thr; 0.66 Met(O₂); 1.50 Arg; 0.68 Pro.

20

25

Example 140

N-Ac-Sar-Gly-Val-D-Ile-Thr-Arg-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Arg(Pmc) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Arg-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 0.54$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1049 (M)⁺; Acid Anal.: 0.92 Sar; 0.74 Gly; 0.86 Val; 2.00 Ile; 0.49 Thr; 2.67 Arg; 1.00 Pro.

Example 141

N-Ac-Sar-Gly-Val-D-Ile-Thr-Tyr-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Tyr(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Tyr-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.048$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1058 (M)⁺; Acid Anal.: 0.88 Sar; 0.99 Gly; 0.97 Val; 1.97 Ile; 0.52 Thr; 0.92 Tyr; 1.58 Arg; 1.08 Pro.

Example 142

N-Ac-Sar-Gly-Val-D-Ile-Thr-Glu-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Glu(OtBu)-OH for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Glu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t =$

2.348 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M)⁺; Acid Anal.: 1.05 Sar; 1.024 Gly; 0.94 Val; 2.67 Ile; 0.47 Thr; 0.94 Glu; 2.20 Arg; 1.09 Pro.

5

Example 143

N-Ac-Sar-Gly-Val-D-Ile-Thr-Lys(Ac)-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Lys(Ac) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Lys(Ac)-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.744 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1065 (M)⁺; Acid Anal.: 1.03 Sar; 0.99 Gly; 0.95 Val; 2.04 Ile; 0.66 Thr; 1.05 Lys; 1.41 Arg; 1.02 Pro.

Example 144

N-Ac-Sar-Gly-Val-D-Ile-Thr-Propargylgly-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Propargylgly for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Propargylgly-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.003 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 990 (M)⁺; Acid Anal.: 1.05 Sar; 1.00 Gly; 0.93 Val; 2.10 Ile; 0.54 Thr; 1.71 Arg; 0.97 Pro.

Example 145

30 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-D-alloIle for Fmoc-D-Ile and Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the

resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.704 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1023 (M)⁺; Acid Anal.: 0.93 Sar; 0.94 Gly; 0.94 Val; 2.10 Ile; 0.51 Thr; 0.87 Glu; 1.45 Arg; 1.03 Pro.

Example 146

N-Ac-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 was used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.685 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1023 (M)⁺; Acid Anal.: 0.98 Sar; 0.74 Gly; 0.95 Val; 1.04 Ile; 0.49 Thr; 1.04 Leu; 0.94 Glu; 1.63 Arg; 0.97 Pro.

Example 147

N-Ac-Bala-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 65 was used but substituting Fmoc-beta-alanine for Fmoc-4-amino-butyric acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Bala-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.92 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1065 (M)⁺; Acid Anal.: 0.99 Sar; 0.99 Gly; 1.00 Val; 1.86Ile; 0.49 Thr; 1.07 Nva; 1.51 Arg; 1.02 Pro.

Example 148

N-Phenylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 60 was used but substituting phenylacetic acid for butyric acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Phenylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.83 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1070 (M)⁺; Acid Anal.: 1.04 Sar; 0.979 Gly; 1.01 Val; 1.90 Ile; 0.59 Thr; 1.09 Nva; 1.53 Arg; 1.03 Pro.

Example 149

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-Azagly-NH₂

To a solution of N-Ac-Sar-Gly-Val-D-Ile-Thr(tBu)-Nva-Ile-Arg(Pmc)-Pro-OH (0.1288 g) in DMF was added semicarbazide hydrochloride (0.222 g) followed by DIEA (0.346 ml) and PyBrop (0.0513 g). The solution was stirred at rt for 36 hr. The solvent was removed in vacuo and the residue was treated with diethyl ether. The solid was filtered and then treated with (9:1) TFA/anisole (3 mL) at rt for 4 hr. The solvent was again removed in vacuo and the residue was treated with diethyl ether. The precipitate was filtered to give the crude product as a solid. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-Azagly-NH₂ as the trifluoroacetate salt; R_t = 2.67 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M)⁺; Acid Anal.: 0.99 Sar; 0.98 Gly; 1.00 Val; 2.13 Ile; 0.56 Thr; 1.09 Nva; 0.92 Arg; 1.02 Pro.

Example 150

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Sar-NHCH₂CH₃

The procedure described in Example 76 was used but substituting Fmoc-Sar-Sieber ethylamide resin for Fmoc-D-Pro-Sieber ethylamide resin. After cleavage of the peptide

from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Sar-NHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.93 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 968 (M)⁺; Acid Anal.: 1.96 Sar; 0.96 Gly; 0.98 Val; 2.07 Ile; 0.55 Thr; 1.05 Nva; 1.49 Arg.

Example 151

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SerNH₂

The procedure described in Example 75 was used but substituting Fmoc-Ser(tBu)-Sieber amide resin for Fmoc-D-Ala-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SerNH₂ as the trifluoroacetate salt: R_t = 2.65 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1053 (M)⁺; Acid Anal.: 0.99 Sar; 0.95 Gly; 1.00 Val; 1.96 Ile; 0.57 Thr; 1.12 Nva; 1.03 Arg; 1.03 Pro; 0.27 Ser.

Example 152

N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 54 was used but substituting Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.85 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1052 (M)⁺; Acid Anal.: 1.01 Sar; 0.93 Gly; 0.95Val; 1.16 Leu; 1.10 Ile; 0.51 Thr; 1.04 Nva; 1.67 Arg; 0.96 Pro.

Example 153

N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Ala for Fmoc-Gly. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.056$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M)⁺; Acid Anal.: 1.32 Sar; 0.96 Ala; 0.94 Val; 2.10 Ile; 0.52 Thr; 0.98 Nva; 1.65 Arg; 1.01 Pro.

Example 154

N-Ac-Sar-Leu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Leu for Fmoc-Gly. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Leu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 3.628$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1050 (M)⁺.

Example 155

N-Ac-Sar-Ser-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Ser(tBu) for Fmoc-Gly. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Ser-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 2.955$ min

(gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M)⁺.

Example 156

5 N-Ac-Sar-Phe-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Phe for Fmoc-Gly. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Phe-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.83 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1084 (M)⁺.

15 Example 157

N-Ac-Sar-Glu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Glu(OtBu)-OH for Fmoc-Gly. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Glu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.08 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1065 (M)⁺.

25 Example 158

N-Ac-Sar-Pro-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Pro for Fmoc-Gly and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions

were lyophilized to yield N-Ac-Sar-Pro-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.343 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1034 (M)⁺.

5

Example 159

N-Ac-Sar-Asn-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Asn(Trt) for Fmoc-Gly and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Asn-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.112 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1051 (M)⁺.

15

Example 160

N-Ac-Sar-Asp-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Asp(OtBu)-OH for Fmoc-Gly and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Asp-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 2.9113 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1052 (M)⁺.

25

Example 161

N-Ac-Asn-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Asn(Trt) for Fmoc-Sar and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude

product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Asn-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.06 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1037 (M)⁺.

Example 162

N-Ac-Gln-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gln(Trt) for Fmoc-Sar and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Gln-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.10 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1051 (M)⁺.

Example 163

N-Ac-Ser-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ser(tBu) for Fmoc-Sar and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Ser-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t = 3.15 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1010 (M)⁺.

Example 164

N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Cit for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting

groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.97 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1080 (M)⁺.

Example 165

N-Ac-Glu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

10 The procedure described in Example 1 was used but substituting Fmoc-Glu(tBu)-OH for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-
15 Ac-Glu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.69 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1052 (M)⁺.

Example 166

20 N-Ac-Gaba-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-gamma-aminobutyric acid for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to
25 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Gaba-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.17 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M)⁺.

Example 167

N-Ac-Bala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-beta-alanine for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Bala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.14 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M)⁺.

Example 168

N-Ac-Gln-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gln(Trt) for Fmoc-Sar. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Gln-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.00 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1051 (M)⁺.

Example 169

N-Ac-Sar-Gly-Gly-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gly for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Gly-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.46 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 952 (M)⁺.

Example 170

N-Ac-Sar-Gly-Glu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Glu(OtBu)-OH for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Glu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 1.74 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M)⁺.

Example 171

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

- The procedure described in Example 4 was used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt: R_t = 2.80 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1037 (M)⁺; Acid Anal.: 0.98 Sar; 0.94 Gly; 0.97 Val; 2.23 Ile; 0.51 Thr; 0.90 Glu; 1.16 Arg; 1.03 Pro.

Example 172

N-Ac-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

- The procedure described in Example 4 was used but substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as

the trifluoroacetate salt: $R_t = 2.90$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1037 (M)⁺; Acid Anal.: 1.05 Sar; 0.97 Gly; 0.99 Val; 1.30 Leu 1.11 Ile; 0.52 Thr; 0.89 Glu; 1.20 Arg; 1.04 Pro.

5 Example 173

H-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 172 was used but omitting the last coupling with acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield H-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt: $R_t = 2.55$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 981 (M)⁺; Acid Anal.: 1.02 Sar; 0.93 Gly; 1.02 Val; 1.05 Leu; 1.02 Ile; 0.55 Thr; 0.84 Gln; 1.31 Arg; 1.03 Pro.

Example 174

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 54 was used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 2.02$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1081 (M)⁺; Acid Anal.: 1.00 Sar; 0.94 Gly; 1.00 Val; 2.00 Ile; 0.52 Thr; 0.87 Gln; 1.37 Arg; 1.05 Pro.

Example 175

30 N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 174 was used but substituting Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the

protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 2.284$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1081 (M)⁺.

Example 176

N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 4 was used but substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Gln(Trt) for Fmoc-Nva. Following the coupling with Fmoc-Sar and protection the resin was treated with succinic anhydride/pyridine as described in Example 54. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt: $R_t = 2.56$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1095 (M)⁺; Acid Anal.: 0.95 Sar; 0.94 Gly; 1.02 Val; 1.02 Leu; 1.05 Ile; 0.56 Thr; 0.86 Gln; 1.00 Arg; 1.07 Pro.

Example 177

N-Ac-Sar-Gly-Val-D-Leu-Thr-Asp-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 146 was used but substituting Fmoc-Asp(OtBu)-OH for Fmoc-Gln(Trt). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Asp-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: $R_t = 2.53$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1010 (M)⁺; Acid Anal.: 1.00 Sar; 0.95 Gly; 1.01 Val; 1.02 Leu; 1.00 Ile; 0.56 Thr; 0.99 Asp; 1.43 Arg; 1.03 Pro.

Example 178

N-Ac-Sar-Gly-Val-D-Ile-Thr-Asp-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 142 was used but substituting Fmoc-
5 Asp(OtBu)-OH for Fmoc-Glu(OtBu)-OH. After cleavage of the peptide from the resin
and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product
was purified by C-18 column chromatography using a solvent mixture varying in a
gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions
were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Asp-Ile-Arg-ProNHCH₂CH₃ as the
10 trifluoroacetate salt; $R_t = 2.455$ min (gradient of 10% to 30% acetonitrile in water
containing 0.01% TFA over 30 min period); MS (ESI) m/e 1010 (M)⁺.

Example 179

N-Ac-Sar-Gly-Val-D-Ile-Thr-Asn-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 43 was used but substituting Fmoc-Asn(Trt)
15 for Fmoc-Gln(Trt). After cleavage of the peptide from the resin and removal of the
protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18
column chromatography using a solvent mixture varying in a gradient of 10% to 50%
acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-
20 Ac-Sar-Gly-Val-D-Ile-Thr-Asn-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; $R_t =$
2.68 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30
min period); MS (ESI) m/e 1009 (M)⁺.

Example 180

N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O)-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 139 was used but substituting Fmoc-Met(O)
25 for Fmoc-Met(O₂). After cleavage of the peptide from the resin and removal of the
protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18
column chromatography using a solvent mixture varying in a gradient of 10% to 50%
30 acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-
Ac-Sar-Gly-Val-D-Ile-Thr-Met(O)-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; R_t

= 2.713 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1042 (M)⁺.

Example 181

5 N-Ac-Sar-Gly-Val-D-Leu-Thr-Asn-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 146 was used but substituting Fmoc-Asn(Trt) for Fmoc-Gln(Trt). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Asn-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.752 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1009 (M)⁺.

15 Example 182

The procedure described in Example 1 is used but separately substituting in the syntheses Fmoc-D-Ile with the following amino acids: Fmoc-D-Thr(tBu), Fmoc-D-Ser(tBu), Fmoc-D-Hser(tBu), Fmoc-D-Gln(Trt), Fmoc-D-Asn(Trt), Fmoc-D-Cit, Fmoc-D-Hcit, Fmoc-D-Hle, Fmoc-D-Neopentylgly. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the trifluoroacetate salt of the following peptides:

N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

25 N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cit-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

30 N-Ac-Sar-Gly-Val-D-Hcit-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Hle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-Neopentylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃.

Example 183

N-Ac-Sar-Gly-Val-D-Ile-Thr-Phe(4-CONH₂)-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 43 is used but substituting Fmoc-Phe[4-CONH(Trt)] for Fmoc-Gln(Trt). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Phe(4-CONH₂)-Ile-Arg-ProNHCH₂CH₃.

Example 184

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-His-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-His(Boc) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-His-ProNHCH₂CH₃.

Example 185

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Isp)-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Lys(N-epsilon-Isp,N-epsilon-Boc) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Isp)-ProNHCH₂CH₃.

Example 186

The procedure described in Example 185 is used but separately substituting in each synthesis Fmoc-Lys(N-epsilon-nicotinyl), Fmoc-Orn(N-delta-nicotinyl), Fmoc-Orn(N-delta-Isp,N-epsilon-Boc), Fmoc-Phe(4-N-Isp,4-N-Boc), Fmoc-Cha(4-N-Isp,4-N-Boc)

instead of Fmoc-Lys(N-epsilon-Isp,N-epsilon-Boc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude products are purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions

are lyophilized to yield the following peptides:

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Nic)-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(Nic)-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(Isp)-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-NIsp)-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cha(4-NIsp)-ProNHCH₂CH₃.

Example 187

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Harg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Harg(Pmc) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Harg-ProNHCH₂CH₃.

Example 188

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Norarg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Norarg(N,N-bis-Boc) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Norarg-ProNHCH₂CH₃.

Example 189

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cit-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Cit for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cit-ProNHCH₂CH₃.

Example 190

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Lys(Boc) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys-ProNHCH₂CH₃.

Example 191

N-Ac-Sar-Gly-Val-D-Ile-Phe(4-CH₂OH)-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Phe[4-CH₂O(Trt)] for Fmoc-Thr(Trt). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Phe(4-CH₂OH)-Nva-Ile-Arg-ProNHCH₂CH₃.

Example 192

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-guanidino)-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Phe(4-bis-Boc-guanidino) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was

purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-guanidino)-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.423 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1042 (M+H)⁺.

Example 193

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Aminopyrimidinylbutanoyl-ProNHCH₂CH₃.

The procedure described in Example 1 was used but substituting Fmoc-2-amino-4-[(2-amino)-pyrimidinyl]butanoic acid for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Aminopyrimidinylbutanoyl-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.303 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1016 (M+H)⁺.

Example 194

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-CH₂NHlsp)-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Phe(4-CH₂Nlsp-Boc) for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-CH₂NHlsp)-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 195

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Gly[4-Pip(N-amidino)]-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Gly-4-piperidinyI[N-amidino(BOC)₂] for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Gly (4-Pip-amidino)-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 196

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala[4-Pip(N-amidino)]-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Ala-[4-piperidinyI-(N',N''-bis-Boc-amidino)] for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala[4-Pip(N-amidino)]-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 197

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-guanidino)-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Ala-[3-(bis-Boc)guanidino] for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-guanidino)-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 198

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-pyrrolidinylamidino)-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Ala[3-pyrroli-dinyl-(2-N,N'-bis-Boc-amidino)] for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala-(3-pyrrolidinyl-amidino)-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 199

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(2-imidazo)-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Orn-[N-2-(1-Boc)imidazoliny] for Fmoc-Arg(Pmc). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(2-imidazo)-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 200

N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 54 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Nva Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 201

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 54 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 202

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva and, after the coupling with Fmoc-Sar, acylating the peptide resin with succinic anhydride as described in Example 54. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

Example 203

N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 201 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 204

N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 202 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

Example 205

N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 175 is used but substituting Fmoc-D-alloIle for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 206

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 205 is used but substituting Fmoc-D-Ile for Fmoc-D-alloIle. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 207

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-D-Ala-NH₂ as the trifluoroacetate salt.

Example 208

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 4 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 209

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

Example 210

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 4 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

5 Example 211

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 209 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

15 Example 212

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 210 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 213

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SarNH₂

25 The procedure described in Example 75 is used but substituting Fmoc-Sar-Seiberamide-resin for Fmoc-D-Ala-Seiberamide-resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SarNH₂ as the trifluoroacetate salt.

Example 214

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-SarNH₂

The procedure described in Example 213 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-SarNH₂ as the trifluoroacetate salt.

Example 215

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-SarNH₂

The procedure described in Example 213 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-SarNH₂ as the trifluoroacetate salt.

Example 216

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-SarNH₂

The procedure described in Example 215 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-SarNH₂ as the trifluoroacetate salt.

Example 217

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 207 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

5 Example 218

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 208 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-
10 water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 219

15 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 15 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-
20 water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 220

N-Ac-Sar-Gly-Val-D-Ile-Thr-Orn(Ac)-Ile-Arg-ProNHCH₂CH₃

25 The procedure described in Example 1 is used but substituting Fmoc-Orn(Ac) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-
30 Val-D-Ile-Thr-Orn(Ac)-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 221

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂

The procedure described in Example 149 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂ as the trifluoroacetate salt.

Example 222

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂

The procedure described in Example 149 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂ as the trifluoroacetate salt.

Example 223

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂

The procedure described in Example 222 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂ as the trifluoroacetate salt.

Example 224

N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 61 is used but substituting tetrahydro-2-furoic acid for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18

column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

5

Example 225

N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 61 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

15

Example 226

N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 225 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

25

Example 227

N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 209 is used but substituting tetrahydro-2-furoic acid for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield

30

N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

Example 228

5 N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 227 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

Example 229

15 N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 4 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-Gln(Trt) for Fmoc-Nva and tetrahydro-2-furoic acid for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 230

25 The procedures described in Examples 224, 225, 226, 227, 228, and 229 are used but substituting N-acetyl-6-aminocaproic acid (6-Ac-Aca) instead of tetrahydro-2-furoyl. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-(6-Ac-Aca)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂, and
 N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 231

The procedures described in Examples 224, 225, 226, 227, 228, and 229 are used but substituting N-acetyl-4-aminobutyric acid (4-Ac-Gaba) instead of N-acetyl-6-aminocaproic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂, and
 N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 232

The procedures described in Examples 224, 225, 226, 227, 228, and 229 are used but substituting 2-furoic acid instead of tetrahydro-2-furoic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,

N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂, and
N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 233

- 5 The procedures described in Examples 224, 225, 226, 227, 228, and 229 are used but substituting shikimic acid instead of tetrahydro-2-furoic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA.
- 10 The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:
- N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
15 N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂, and
N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 234

- 20 The procedures described in Examples 224, 225, 226, 227, 228, and 229 are used but substituting 2-methyl-nicotinic acid instead of tetrahydro-2-furoic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides:
- 25 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂, and
30 N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 235

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile and Fmoc-Leu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

Example 236

N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 4 is used but substituting Fmoc-Leu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 237

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 73 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 238

N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-Leu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting

groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

5

Example 239

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-Leu for Fmoc-Nva and acylating with succinic anhydride after the coupling with Fmoc-Sar and deprotection as described in Example 54. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

15

Example 240

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 206 is used but substituting Fmoc-Leu for Fmoc-Gln(Trt) and acylating with succinic anhydride after the coupling with Fmoc-Sar and deprotection as described in Example 54. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

25

Example 241

The procedures described in Examples 201, 202 and 203 are used but substituting Fmoc-Leu instead of Fmoc-Gln(Trt). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of

30

10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides:

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃, and

5 N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂

Example 242

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-AzaglyNH₂

The procedure described in Example 149 is used but substituting Fmoc-Leu for
10 Fmoc-Nva and acylating with succinic anhydride after the coupling with Fmoc-Sar and deprotection as described in Example 54. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized
15 to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-AzaglyNH₂ as the trifluoroacetate salt.

Example 243

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHethyl-(1-pyrrolidine)

20 The procedure described in Example 5 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-
25 Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHethyl-(1-pyrrolidine) as the trifluoroacetate salt.

Example 244

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNH(ethyl-1-cyclohexyl)

The procedure described in Example 8 is used but substituting Fmoc-D-alloIle for
30 Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-

water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNH(ethyl-1-cyclohexyl) as the trifluoroacetate salt.

Example 245

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHethyl-(1-pyrrolidine)

The procedure described in Example 5 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHethyl-(1-pyrrolidine) as the trifluoroacetate salt.

Example 246

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl)

The procedure described in Example 8 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl) as the trifluoroacetate salt.

Example 247

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl)

The procedure described in Example 246 is used but acylating the peptide resin with succinic anhydride after the coupling with Fmoc-Sar and deprotection as described in Example 54. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl) as the trifluoroacetate salt.

Example 248

The procedures described in Examples 11 is used but substituting the appropriate protected amino acids as described in Examples 14, 43, 74, 73, 54, 174, and 132 respectively. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 10 N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 15 N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃, and
 N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃.

Example 249

20 The procedures described in Examples 49 is used but substituting the appropriate protected amino acids as described in Examples 14, 4, 75, 54 and 132 respectively. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01%
 25 TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Allygly-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Ac-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 30 N-Ac-Sar-Gly-Val-D-alloIle-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 N-Ac-Sar-Gly-Val-D-Ile-Ser-Allygly-Ile-Arg-Pro-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-Leu-Ser-Allygly-Ile-Arg-Pro-ProNHCH₂CH₃

Example 250

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-SerNH₂

- 5 The procedure described in Example 75 is used but substituting Fmoc-D-Ser(tBu)-Sieber amide resin for Fmoc-D-Ala-Sieber amide resin. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-SerNH₂ as the trifluoroacetate salt.
- 10

Example 251

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHOH

- 15 The procedure described in Example 149 is used but hydroxylamine hydrochloride for semicarbazide hydrochloride. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHOH as the trifluoroacetate salt.
- 20

Example 252

N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

- 25 The procedure described in Example 132 is used but substituting Fmoc-D-Ile for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

30

Example 253

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 132 is used but substituting Fmoc-D-alloIle for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 254

N-Ac-Sar-Gly-Val-D-Leu-Hser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 132 is used but substituting Fmoc-Hser(tBu) for Fmoc-Ser(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Hser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 255

N-Ac-Sar-Gly-Gln-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Gln(Trt) for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Gln-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt; *R*_t = 2.36 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) *m/e* 1023 (M)⁺.

Example 256

N-Ac-Sar-Gly-Nva-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Nva for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Nva-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.28 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 994 (M)⁺.

Example 257

N-Ac-Sar-Gly-Ile-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Ile for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Ile-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.55 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M)⁺.

Example 258

N-Ac-Sar-Gly-Phe-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 was used but substituting Fmoc-Phe for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Phe-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.77 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1042 (M)⁺.

Example 259

N-Ac-Sar-Gly-Leu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- The procedure described in Example 1 was used but substituting Fmoc-Leu for
- 5 Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Leu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 3.56 min
- 10 (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1008 (M)⁺.

Example 260

N-Ac-Sar-Gly-Ser-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- 15 The procedure described in Example 1 was used but substituting Fmoc-Ser(tBu) for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-
- 20 Ser-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt: R_t = 2.41 min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 982 (M)⁺.

Example 261

N-Ac-Thr-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

- 25 The procedure described in Example 1 was used but substituting Fmoc-Thr(tBu) for Fmoc-Sar and Fmoc-D-Leu for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product was purified by C-18 column chromatography using a solvent mixture varying in
- 30 a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Thr-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as

the trifluoroacetate salt: $R_t = 3.33$ min (gradient of 10% to 30% acetonitrile in water containing 0.01% TFA over 30 min period); MS (ESI) m/e 1024 (M)⁺.

Example 262

The procedures described in Example 46 is used but substituting the appropriate protected amino acids as describes in Examples 75, 4, 54, and 132. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA.

The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ala-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,

N-Ac-Sar-Gly-Val-D-Ile-Ser-Ala-Ile-Arg-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-Leu-Ser-Ala-Ile-Arg-ProNHCH₂CH₃.

Example 263

The procedures described in Example 262 is used but substituting Fmoc-Val for Fmoc-Ala. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Val-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,

N-Ac-Sar-Gly-Val-D-Ile-Ser-Val-Ile-Arg-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-Leu-Ser-Val-Ile-Arg-ProNHCH₂CH₃.

Example 264

The procedures described in Example 263 is used but substituting Fmoc-DNva for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

- 10 N-Ac-Sar-Gly-Val-D-alloIle-Thr-D-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
N-Ac-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-Val-D-alloIle-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
N-Succinyl-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
15 N-Ac-Sar-Gly-Val-D-Ile-Ser-D-Nva-Ile-Arg-Pro-ProNHCH₂CH₃, and
N-Ac-Sar-Gly-Val-D-Leu-Ser-D-Nva-Ile-Arg-Pro-ProNHCH₂CH₃.

Example 265

N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

20 The procedure described in Example 132 is used but substituting Fmoc-D-Ile for Fmoc-D-Leu and Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are
25 lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 266

N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

30 The procedure described in Example 132 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

5

Example 267

N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 75 is used but substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

15

Example 268

N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 267 is used but substituting Fmoc-D-Ile for Fmoc-D-Leu and Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate salt.

25

Example 269

N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 54 is used but substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are

30

lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 270

5 N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 269 is used but substituting Fmoc-D-Ile for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-
10 water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 271

15 N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 270 is used but substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are
20 lyophilized to yield N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 272

25 N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 270 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-
30 Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 273

N-Ac-Sar-Gly-Val-D-Ile-Ser-Ser-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 265 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Ser-Ser-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 274

N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 266 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 275

N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 13 is used but substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

Example 276

N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 13 is used but substituting Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the

protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate salt.

5

Example 277

N-Ac-Sar-Gly-Val-D-Leu-Ser-Leu-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 132 is used but substituting Fmoc-Leu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Leu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

10

Example 278

N-Ac-Sar-Gly-Val-D-Ile-Ser-Leu-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 277 is used but substituting Fmoc-D-Ile for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Ser-Leu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

15

20

Example 279

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 132 is used but substituting Fmoc-D-alloIle for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

25

30

Example 280

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 265 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 281

N-Succinyl-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 270 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Succinyl-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 282

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂

The procedure described in Example 276 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂ as the trifluoroacetate.

Example 283

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂

The procedure described in Example 268 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18

column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂ as the trifluoroacetate.

5

Example 284

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Leu-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 265 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile and Fmoc-Leu for Fmoc-Gln(Trt). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Ser-Leu-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

10

15

Example 285

N-Ac-Sar-Gly-Val-D-alloIle-Ser-Ser-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 276 is used but substituting Fmoc-D-alloIle for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloIle-Ser-Ser-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

20

25

Example 286

The procedure described in Example 125 is used but separately substituting Fmoc-D-Ile and Fmoc-D-alloIle, respectively, for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to the the following peptides as trifluoroacetate salt:

30

N-Ac-Sar-Gly-Val-D-Ile-Gly-Nva-Ile-Arg-ProNHCH₂CH₃ and

N-Ac-Sar-Gly-Val-D-alloIle-Gly-Nva-Ile-Arg-ProNHCH₂CH₃.

Example 287

The procedure described in Example 125 and 286 is used but separately substituting Fmoc-D-Ile and Fmoc-D-alloIle, respectively, for Fmoc-D-Leu and substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the trifluoroacetate salt of:

N-Ac-Sar-Gly-Val-D-Leu-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Gly-Gln-Ile-Arg-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-alloIle-Gly-Gln-Ile-Arg-ProNHCH₂CH₃.

Example 288

The procedure described in Example 123 is used but separately substituting Fmoc-D-Ile and Fmoc-D-alloIle, respectively, for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the trifluoroacetate salt of:

N-Ac-Sar-Gly-Val-D-Ile-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃ and

N-Ac-Sar-Gly-Val-D-alloIle-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃.

Example 289

The procedure described in Example 123 and 288 is used but separately substituting Fmoc-D-Ile and Fmoc-D-alloIle, respectively, for Fmoc-D-Leu and substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the trifluoroacetate salt of:

N-Ac-Sar-Gly-Val-D-Leu-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃, and

N-Ac-Sar-Gly-Val-D-alloIle-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃.

Example 290

N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Ser(tBu) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 291

N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Thr(tBu) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 292

N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Gln(Trt) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 293

N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Asn(Trt) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 294

N-Ac-Sar-Gly-Val-D-Arg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Arg(Pmc) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Arg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 295

N-Ac-Sar-Gly-Val-D-3-Pal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-3-Pal for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-3-Pal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

5

Example 296

N-Ac-Sar-Gly-Val-D-Glu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Glu(OtBu)-OH for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Glu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

10

Example 297

N-Ac-Sar-Gly-Val-D-Asp-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Asp(OtBu)-OH for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Asp-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

20

Example 298

N-Ac-Sar-Gly-Val-D-His-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-His(Boc)-OH for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-His-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

25

30

Example 299

N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Hser(tBu) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 300

N-Ac-Sar-Gly-Val-D-alloThr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-alloThr(tBu) for Fmoc-D-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloThr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 301

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-D-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-D-Ile for Fmoc-Ile. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-D-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 302

N-Ac-Sar-Gly-Val-D-Ser-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 290 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Ser-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

5

Example 303

N-Ac-Sar-Gly-Val-D-Thr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 291 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Thr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

10

Example 304

N-Ac-Sar-Gly-Val-D-alloThr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 300 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloThr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

15

20

Example 305

N-Ac-Sar-Gly-Val-D-Ser-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 290 is used but substituting Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to N-Ac-Sar-Gly-Val-D-Ser-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

25

30

Example 306

N-Ac-Sar-Gly-Val-D-Thr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 291 is used but substituting Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Thr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 307

N-Ac-Sar-Gly-Val-D-alloThr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 300 is used but substituting Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloThr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 308

N-Ac-Sar-Gly-Val-D-alloThr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 304 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-alloThr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 309

N-Ac-Sar-Gly-Val-D-Thr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 303 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column

chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Thr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

5

Example 310

The procedure described in Examples 132 and 266 is used but substituting N-acetyl-6-aminocaproic acid (6-Ac-Aca) for acetic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂, and

N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂.

15

Example 311

The procedure described in Examples 310 is used but substituting N-acetyl-gamma-aminobutyric acid (4-Ac-Gaba) instead of N-acetyl-6-aminocaproic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:


N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂, and

N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂.

25


Example 312

The procedure described in Examples 311 is used but substituting 2-furoic acid instead of N-acetyl-gamma-aminobutyric acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:




Example 313

The procedure described in Examples 311 is used but substituting shikimic acid instead of 2-furoic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:




Example 314

The procedure described in Examples 311 is substituting shikimic acid instead of 2-furoic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:




Example 315

The procedure described in Examples 312 is used but substituting 2-methyl-nicotinic acid instead of 2-furoic acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:




Example 316



The procedure described in Example 8 is used but substituting Fmoc-DLeu for Fmoc-DIle and Fmoc-Ser(tBu) for Fmoc-Thr(tBu). After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate.

Example 317



The procedure described in Example 8 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-DIle-Thr-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate.

Example 318

N-Ac-Sar-Gly-Val-DIle-Thr-Leu-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl

The procedure described in Example 8 is used but substituting Fmoc-Leu for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-DIle-Thr-Leu-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate.

Example 319

N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl

The procedure described in Example 8 is used but substituting Fmoc-D-Leu for Fmoc-DIle. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate.

Example 320

N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl

The procedure described in Example 316 is used but substituting Fmoc-Ser(tBu) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate.

Example 321

N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl

The procedure described in Example 316 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl as the trifluoroacetate.

Example 322

N-Ac-Sar-Gly-Val-DIle-Thr-Nva-Ile-Arg-ProNHethyl-1-(S)-cyclohexyl

The procedure described in Example 8 is used but substituting (S)-1-cycloxyethylamine for (R)-1-cycloxyethylamine. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-DIle-Thr-Nva-Ile-Arg-ProNHethyl-1-(S)-cyclohexyl as the trifluoroacetate salt.

Example 323

The procedures described in Example 98 is used but substituting the appropriate protected amino acids as describes in Examples 132, 43, 54, and 75. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA.

The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Pen-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Pen-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 5 N-Ac-Sar-Gly-Val-D-Pen-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Ac-Sar-Gly-Val-D-Pen-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 10 N-Ac-Sar-Gly-Val-D-Pen-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Leu-Ile-Arg-ProNHCH₂CH₃, and
 N-Succinyl-Sar-Gly-Val-D-Pen-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 324

N-Ac-Sar-Gly-Val-D-Cys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 98 is used but substituting Fmoc-D-Cys(Trt)
 for Fmoc-D-Pen(Trt). After cleavage of the peptide from the resin and removal of the
 protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18
 20 column chromatography using a solvent mixture varying in a gradient of 10% to 50%
 acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield N-
 N-Ac-Sar-Gly-Val-D-Cys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate.

Example 325

The procedures described in Example 324 is used but substituting the appropriate
 protected amino acids as describes in Examples 132, 43, 54, and 75. After cleavage of the
 peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3

mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA.

The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

5 N-Ac-Sar-Gly-Val-D-Cys-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,

10 N-Ac-Sar-Gly-Val-D-Cys-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Cys-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,

15 N-Ac-Sar-Gly-Val-D-Cys-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Ser-Ile-Arg-ProNHCH₂CH₃, and

N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Leu-Ile-Arg-ProNHCH₂CH₃.

Example 326

20 N-Ac-Sar-Gly-Pen-DIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Pen(Trt) for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA.

25 The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Pen-DIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 327

N-Ac-Sar-Gly-Cys-DIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 1 is used but substituting Fmoc-Cys(Trt) for Fmoc-Val. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Cys-DIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 328

The procedures described in Example 326 is used but substituting the appropriate protected amino acids as describes in Examples 14, 15, 132, 43, 54, and 75. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Pen-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Pen-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Pen-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,

N-Succinyl-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃, and

N-Succinyl-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 329

N-Ac-Sar-Gly-Val-D-Leu-Pen-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 120 is used but substituting Fmoc-Pen(Trt) for Fmoc-Ala. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Leu-Pen-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 330

The procedures described in Example 329 is used but substituting the appropriate protected amino acids as describes in Examples 14, 15, 132, 43, 54, and 75. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-alloIle-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Pen-Ser-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Pen-Leu-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-Pro-D-AlaNH₂,

N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂CH₃, and

N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂(CH₃)₂.

Example 331

N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 11 is used but substituting Fmoc-Pen(Trt) for

- 5 Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 332

The procedures described in Example 331 is used but substituting the appropriate protected amino acids as describes in Examples 14, 15, 132, 43, 54, and 75. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1)

- 15 TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,

- 20 N-Ac-Sar-Gly-Val-D-Leu-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-Pro-D-AlaNH₂,

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂(CH₃)₂,

N-Ac-Sar-Gly-Val-D-Leu-Ser-Pen-Ile-Arg-ProNHCH₂CH₃,

- 25 N-Ac-Sar-Gly-Val-D-Leu-Gly-Pen-Ile-Arg-ProNHCH₂CH₃, and

N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Pen-Ile-Arg-ProNHCH₂CH₃.

Example 333

N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 96 is used but substituting Fmoc-Gln(Trt) for Fmoc-Nva. After cleavage of the peptide from the resin and removal of the protecting groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

Example 334

The procedures described in Example 333 is used but substituting the appropriate protected amino acids as describes in Examples 132, 43, 54, and 75. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Gln-Ile-Arg-ProNHCH₂CH₃, and
N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Ser-Ile-Arg-ProNHCH₂CH₃.

Example 335

N-Ac-Sar-Ala-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in Example 153 is used but substituting Fmoc-DalloIle for Fmoc-DIle. After cleavage of the peptide from the resin and removal of the protecting

groups the crude product was purified by C-18 column chromatography using solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Ala-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as the trifluoroacetate salt.

5

Example 336

The procedures described in Example 335 is used but substituting the appropriate protected amino acids as described in Examples 132, 43, 54, and 75. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-Ac-Sar-Ala-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Ala-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Ala-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Ac-Sar-Ala-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-ProNHCH₂CH₃,

N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-ProNHCH₂(CH₃)₂, and

N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-Pro-D-AlaNH₂.

Example 337

The procedure described in Example 231 used but substituting N-acetyl-beta-alanine (3-Ac-Bala) for N-acetyl-4-aminobutyric acid. After cleavage of the peptide from the resin and removal of the protecting groups using (9:1) TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01% TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate salt:

N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂.

N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-DAlaNH₂,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 N-(3-Ac-Bala)-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Ala-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Ala-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 N-(3-Ac-Bala)-Sar-Ala-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃, and
 10 N-(3-Ac-Bala)-Sar-Ala-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃.

Example 338

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH

The procedure described in Example 1 is used but substituting omitting the
 15 coupling with ethylamine. After cleavage of the peptide from the resin and removal of the
 protecting groups the crude product was purified by C-18 column chromatography using
 solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01%
 TFA. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-
 Arg-Pro-OH as the trifluoroacetate salt.

Example 339

The procedures described in Example 338 is used but substituting the appropriate
 protected amino acids as described in Examples 14, 15, 132, 43, 54, and 75. After
 cleavage of the peptide from the resin and removal of the protecting groups using (9:1)
 25 TFA/anisole (3 mL) the crude product is purified by C-18 column chromatography using a
 solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.01%
 TFA. The pure fractions are lyophilized to yield the following peptides as trifluoroacetate
 salt:

N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-OH,

30 N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-Pro-OH,

N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-Pro-OH,

N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-Pro-OH,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-OH,
N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-OH,
N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-Pro-OH,
5 N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH, and
N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-Pro-OH.

Example 340

N-Ac-Sar-Gly-Asp-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

10 In an Applied Biosystems 433A peptide synthesizer, 0.1 mM of Fmoc-Pro-Sieber
ethylamide resin was placed in the reaction vessel and cartridges of 1 mM amino acids
were sequentially loaded. The Fastmoc 0.1 with previous peak monitoring protocol was
used. The following is the synthetic cycle:

1. Solvating resin with N-methylpyrrolidone (NMP) for about 5 minutes;
- 15 2. Washing with NMP for about 5 minutes;
3. Removing the Fmoc group using 50% piperidine solution in NMP for 5 minutes,
washing and repeating the process 3 to 4 times;
4. Activating the Fmoc-amino acid using 1 mM of 0.5 M solution of 2-(1H-
benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU)
20 in DMF;
5. Adding the activated Fmoc-amino acid to the reaction vessel followed by 1 mM
of 2 M diisopropylamine in NMP solution;
6. Coupling of the Fmoc-amino acid for 20 minutes;
7. Washing and removal of Fmoc-group using 50 % piperidine in NMP.

25 The following protected amino acids were sequentially coupled to the resin using above
protocol in Table 2, below:

Table 2

Amino acid	Coupling time
1. Fmoc-Arg(Pmc)	20 minutes
2. Fmoc-Ile	20 minutes
3. Fmoc-Nva	20 minutes
4. Fmoc-Thr(<i>t</i> -Bu)	20 minutes
5. Fmoc-D-Leu	20 minutes
6. Fmoc-Asp(OtBu)	20 minutes
7. Fmoc-Gly	20 minutes
8. Fmoc-Sar	20 minutes
9. acetic acid	20 minutes

Upon completion of the synthesis the resin-bound peptide was washed with methanol three times and dried *in vacuo*, then treated with a (95:5) TFA/water solution (3mL) at room temperature overnight. The resin was filtered and washed 3 times with methanol. The filtrates and the washes were combined and concentrated *in vacuo*. The residue was treated with ether and the precipitate was filtered to give the crude peptide as amorphous powder. This was purified by preparative HPLC using C-18 column with a mixture of solvents varying in a gradient from 5 % to 100 % acetonitrile/water containing 0.1% TFA over a period of 50 min. The pure fractions were lyophilized to yield N-Ac-Sar-Gly-Asp-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as trifluoroacetate salt; $R_t = 2.23$ min (using C-18 column and solvents mixture varying in a gradient from 20 % to 95 % acetonitrile/water containing 10 mM ammonium acetate over a period of 10 min); MS (ESI) m/e 1010 (M^+).

Example 341

N-Ac-Sar-Gly-Ala-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in example 340 was used but substituting Fmoc-Ala for Fmoc-Asp(OtBu). After cleavage of the peptide from the resin and removal of the protecting groups the product was precipitated with ether. The crude compound was purified by preparative HPLC to give N-Ac-Sar-Gly-Ala-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as trifluoroacetate salt; $R_t = 2.828$ min (using C-18 column and solvents

mixture varying in a gradient from 20 % to 95 % acetonitrile/water containing 10 mM ammonium acetate over a period of 10 min); MS (ESI) m/e 966 (M^+).

Example 342

5 N-Ac-Sar-Gly-Cha-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in example 340 was used but substituting Fmoc-Cha for Fmoc-Asp(OtBu). After cleavage of the peptide from the resin and removal of the protecting groups the product was precipitated with ether. The crude compound was purified by preparative HPLC to give N-Ac-Sar-Gly-Cha-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as trifluoroacetate salt; R_t = 4.48 min (using C-18 column and solvents mixture varying in a gradient from 20 % to 95 % acetonitrile/water containing 10 mM ammonium acetate over a period of 10 min); MS (ESI) m/e 1048 (M^+).

Example 343

15 N-Ac-Sar-Gly-Met-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in example 340 was used but substituting Fmoc-Met for Fmoc-Asp(OtBu) and Fmoc-D-Ile for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups the product was precipitated with ether. The crude compound was purified by preparative HPLC to give N-Ac-Sar-Gly-Met-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as trifluoroacetate salt; R_t = 3.25 min (using C-18 column and solvents mixture varying in a gradient from 20 % to 95 % acetonitrile/water containing 10 mM ammonium acetate over a period of 10 min); MS (ESI) m/e 1026 ($M+H$); Amino Acid Anal.: 1.09 Sar; 0.97 Gly; 0.94 Met; 2.08 Ile; 0.47 Thr; 1.00 Nva; 1.34 Arg; 1.01 Pro.

Example 344

25 N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in example 340 was used but substituting Fmoc-Cit for Fmoc-Sar, Fmoc-Val for Fmoc-Asp(OtBu) and Fmoc-D-Ile for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups the product was precipitated with ether. The crude compound was purified by preparative HPLC to

N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃ as trifluoroacetate salt; R_t = 2.97 min (using C-18 column and solvents mixture varying in a gradient from 20 % to 95 % acetonitrile/water containing 10 mM ammonium acetate over a period of 10 min); MS (ESI) m/e 1080 (M+H); Amino Acid Anal.: 0.98 Cit; 0.93 Gly; 0.98 Val; 2.05 Ile; 0.51 Thr; 0.99 Nva; 1.37 Arg; 1.01 Pro.

Example 345

N-Ac-Sar-Gly-Val-D-Ile-Thr-Hser-Ile-Arg-ProNHCH₂CH₃

The procedure described in example 340 was used but substituting Fmoc-Hser(t-Bu) for Fmoc-Nva, Fmoc-Val for Fmoc-Asp(OtBu), and Fmoc-D-Ile for Fmoc-D-Leu. After cleavage of the peptide from the resin and removal of the protecting groups the product was precipitated with ether. The crude compound was purified by preparative HPLC to give N-Ac-Sar-Gly-Val-D-Ile-Thr-Hser-Ile-Arg-ProNHCH₂CH₃ as trifluoroacetate salt; R_t = 2.782 min (using C-18 column and solvents mixture varying in a gradient from 10 % to 40 % acetonitrile/water containing 0.1 % TFA over a period of 33 min); MS (ESI) m/e 996 (M+H); Amino Acid Anal.: 1.00 Sar; 0.95 Gly; 1.01 Val; 1.99 Ile; 0.60 Thr; 1.07 Arg; 1.04 Pro.

Example 346

N-Ac-Sar-Gly-Val-D-alloIle-His-Nva-Ile-Arg-ProNHCH₂CH₃

The procedure described in example 340 was used but substituting Fmoc-Val for Fmoc-Asp(OtBu), Fmoc-D-alloIle for Fmoc-D-Leu, and Fmoc-His(Boc) for Fmoc-Thr(t-Bu). After cleavage of the peptide from the resin and removal of the protecting groups the product was precipitated with ether. The crude compound was purified by preparative HPLC to give N-Ac-Sar-Gly-Val-DalloIle-His-Nva-Ile-Arg-ProNHCH₂CH₃ as a trifluoroacetate salt; R_t = 3.12 min (using C-18 column and solvents mixture varying in a gradient from 10 % to 40 % acetonitrile/water containing 0.1 % TFA over a period of 33 min); MS (ESI) m/e 1030 (M⁺).

Example 347

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-Butyl

Resin Preparation

- 5 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin (0.5 g, 0.54 mmol/g substitution) was placed in a solid phase synthesis reaction vessel containing (9:1) DMA/acetic acid (4 mL). The mixture was shaken for 5 min. The resin was drained and this process was repeated three times. To the swollen resin were added 10-15 grains of activated 4A molecular sieves and (9:1) DMA/acetic acid (4mL) and 10 molar equivalents
- 10 of n-butylamine. The slurry was shaken for 1 h at rt and to it was added 10 molar equivalents of sodium triacetoxyborohydride. The slurry was shaken for 2 h at rt. The resin was drained and washed three times with DMA, three times with methanol, three times with dichloromethane, three times with diethyl ether and dried *in vacuo* at rt overnight. The dry resin was swollen in DMA (4 mL) and shaken for 5 min. This process
- 15 was repeated twice.

Coupling of Fmoc-Pro

- To the swollen resin in the reaction vessel were added sequentially the following chemicals: DMA (4 mL), one equivalent of DIEA, a DMA solution containing 3.0 equivalents of Fmoc-Pro, 3.0 equivalents of HATU, and 3.0 equivalents of DIEA. The
- 20 slurry was shaken overnight. The resin was drained and washed three times with DMA, three times with methanol, three times with dichloromethane, three times with diethyl ether and dried *in vacuo* at rt overnight. A small portion of the resin was used to determine the Fmoc-Pro loading. The rest of the resin was shaken with DMA (4 mL) three times for 5 min and then for 1 h at rt with a solution of (8:1:1) DMA/pyridine/acetic
- 25 anhydride (5 mL). The resin was drained and washed three times with DMA, three times with methanol, three times with dichloromethane, and three times with diethyl ether. The resin was dried *in vacuo* at rt overnight and then used in the subsequent solid phase peptide synthesis.

Synthesis of Peptide

- 30 In the synthesis of the above peptide, the amino acids, the coupling conditions and the synthetic protocol used were identical to those described in Example 340. Upon

completion of the synthesis the peptide and the protecting groups were cleaved at rt using (95:5) TFA/water (3 mL) for 3h. The resin was filtered and washed three times with methanol. The combined filtrates were concentrated *in vacuo* and to the residue was added diethylether. The solid precipitate was filtered. The crude product was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to yield NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-Butyl as the trifluoroacetate salt: $R_t = 3.792$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1022 (M^+).

Example 348

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Butyl

The procedure described in example 347 was used but substituting isobutylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Butyl as the trifluoroacetate salt: $R_t = 3.731$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1022 (M^+).

Example 349

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Amyl

The procedure described in example 347 was used but substituting isoamylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Amyl as the trifluoroacetate salt: $R_t = 4.086$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1036 (M^+).

Example 350

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-hexyl

The procedure described in example 347 was used but substituting in n-hexylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-Hexyl as the trifluoroacetate salt: $R_t = 4.527\text{min}$ (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1050 (M^+).

Example 351

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3,3-dimethyl)butyl

The procedure described in example 347 was used but substituting (3,3-dimethyl)butylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3,3-dimethyl)butyl as the trifluoroacetate salt: $R_t = 4.366\text{min}$ (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1050 (M^+).

Example 352

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-ethoxy)ethyl

The procedure described in example 347 was used but substituting (2-ethoxy)ethylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-ethoxy)ethyl as the trifluoroacetate salt: $R_t =$

3.356 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1038 (M⁺).

Example 353

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-isopropoxy)ethyl

The procedure described in example 347 was used but substituting (2-isopropoxy)ethylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg- ProNH-(2-isopropoxy)ethyl as the trifluoroacetate salt: R_t = 3.57 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1052 (M⁺).

Example 354

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3-methoxy)propyl

The procedure described in example 347 was used but substituting (3-methoxy)propylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg- ProNH-(3-methoxy)propyl as the trifluoroacetate salt: R_t = 3.26 min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH₄Ac over 10 min period); MS (ESI) m/e 1038 (M⁺).

Example 355

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(cyclopentyl)methyl

The procedure described in example 347 was used but substituting (cyclopentyl)methylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg- ProNH-(cyclopentyl)methyl as the trifluoroacetate salt: $R_t = 4.148$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1048 (M^+).

Example 356

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-cyclohexyl

The procedure described in example 347 was used but substituting cyclohexylamine for n-butylamine. After cleavage of the peptide from the resin and precipitation with ether the crude product was obtained. This was purified by C-18 column chromatography using a solvent mixture varying in a gradient of 10% to 50% acetonitrile-water containing 0.1% TFA. The pure fractions were lyophilized to NAcSar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg- ProNH-cyclohexyl as the trifluoroacetate salt: $R_t = 4.070$ min (gradient of 20% to 95% acetonitrile in water containing 0.01 M NH_4Ac over 10 min period); MS (ESI) m/e 1048 (M^+).

In Vitro Assay for Angiogenic Activity

The human microvascular endothelial (HMVEC) migration assay was run according to the procedure of S. S. Tolsma, O. V. Volpert, D. J. Good, W. F. Frazier, P. J. Polverini and N. Bouck, J. Cell Biol. 122, 497-511 (1993).

The HMVEC migration assay was carried out using Human Microvascular Endothelial Cells-Dermal (single donor) and Human Microvascular Endothelial Cells, (neonatal). The BCE or HMVEC cells were starved overnight in DME containing 0.1% bovine serum albumin (BSA). Cells were then harvested with trypsin and resuspended in DME with 0.1% BSA at a concentration of 1.5×10^6 cells per ml. Cells were added to

the bottom of a 48 well modified Boyden chamber (Nucleopore Corporation, Cabin John, MD). The chamber was assembled and inverted, and cells were allowed to attach for 2 hours at 37 °C to polycarbonate chemotaxis membranes (5 µm pore size) that had been soaked in 0.1% gelatin overnight and dried. The chamber was then reinverted, and test substances (total volume of 50 µl), including activators, 15 ng/ml bFGF/VEGF, were added to the wells of the upper chamber. The apparatus was incubated for 4 hours at 37 °C. Membranes were recovered, fixed and stained (Diff Quick, Fisher Scientific) and the number of cells that had migrated to the upper chamber per 3 high power fields counted. Background migration to DME + 0.1 BSA was subtracted and the data reported as the number of cells migrated per 10 high power fields (400X) or, when results from multiple experiments were combined, as the percent inhibition of migration compared to a positive control.

The compounds described in Examples 1 to 339 inhibited human endothelial cell migration in the above assay from about 30% to about 95% inhibition when tested at concentrations of 10 nM or 20 nM, as reported below in Table 3.

Table 3.
In Vitro Angiogenic Activity


<u>Ex. #</u>	<u>% Inhib. @ 20 nM</u>	<u>% Inhib. @ 10 nM</u>
1	87.3	76.9
3	56.0	---
4	71.3	---
5	---	87.2
8	---	88.2
11	70.4	---
12	55.8	---
18	---	51.4
28	---	47.0
42	60.2	---
43	---	94.1
46	77.5	---
47	69.7	---
49	83.4	---

<u>Ex. #</u>	<u>% Inhib. @ 20 nM</u>	<u>% Inhib. @ 10 nM</u>
50	71.6	---
51	67.0	---
52	46.5	---
53	76.7	---
54	81.3	---
55	59.2	---
56	49.9	---
57	56.6	---
58	68.8	---
59	82.3	---
60	75.3	---
61	---	83.7
63	---	82.4
66	76.1	---

5

WHAT IS CLAIMED IS:

1. A compound of the formula:



- 5 or a pharmaceutically acceptable salt, ester, solvate or prodrug thereof, wherein:

A_0 is hydrogen or an acyl group selected from:

- (1) $R-(CH_2)_n-C(O)-$; wherein n is an integer from 0 to 8 and R is selected from hydroxyl; methyl; N-acetylamino; methoxyl; carboxyl; cyclohexyl optionally containing a one or two double bonds and optionally substituted with one to three hydroxyl groups; and a 5- or 6-membered ring aromatic or nonaromatic ring optionally containing one or two heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the ring is optionally substituted with a moiety selected from alkyl, alkoxy, and halogen; and
- (2) $R^1-CH_2CH_2-(OCH_2CH_2O)_p-CH_2-C(O)-$; wherein R^1 is selected from hydrogen, alkyl, and N-acetylamino, and p is an integer from 1 to 8;

A_1 is an amino acyl residue selected from:

- (1) alanyl,
(2) asparaginyI,
(3) citrullyl,
(4) glutaminyI,
(5) glutamyl,
(6) N-ethylglycyl,
(7) methionyl,
(8) N-methylalanyl,
(9) prolyl,
(10) pyro-glutamyl,
(11) sarcosyl,

- 30 (12) seryl,
 (13) threonyl,
 (14) $-\text{HN}-(\text{CH}_2)_q-\text{C}(\text{O})-$, wherein q is 1 to 8, and
 (15) $-\text{HN}-\text{CH}_2\text{CH}_2-(\text{OCH}_2\text{CH}_2\text{O})_r-\text{CH}_2-\text{C}(\text{O})-$, wherein r is 1 to 8;

A₂ is an amino acyl residue selected from:

- 35 (1) alanyl,
 (2) asparaginy, l,
 (3) aspartyl,
 (4) glutaminy, l,
 (5) glutamyl,
 40 (6) leucyl,
 (7) methionyl,
 (8) phenylalanyl,
 (9) prolyl,
 (10) seryl,
 45 (11) $-\text{HN}-(\text{CH}_2)_q-\text{C}(\text{O})-$, wherein q is 1 to 8, and
 (12) $-\text{HN}-\text{CH}_2\text{CH}_2-(\text{OCH}_2\text{CH}_2\text{O})_r-\text{CH}_2-\text{C}(\text{O})-$, wherein r is 1 to 8;

A₃ is an amino acyl residue selected from:

- (1) alanyl,
 (2) asparaginy, l,
 50 (3) citrullyl,
 (4) cyclohexylalanyl,
 (5) cyclohexylglycyl,
 (6) glutaminy, l,
 (7) glutamyl,
 55 (8) glycyl,
 (9) isoleucyl,
 (10) leucyl,
 (11) methionyl,
 (12) norvalyl,
 60 (13) phenylalanyl,
 (14) seryl,

- 65
- (15) *t*-butylglycyl,
 - (16) threonyl,
 - (17) valyl,
 - (18) penicillaminyl, and
 - (19) cystyl;

A₄ is an amino acyl residue of L or D configuration selected from:

- 70
- (1) allo-isoleucyl,
 - (2) glycyl,
 - (3) isoleucyl,
 - (4) prolyl,
 - (5) dehydroleucyl,
 - (6) D-alanyl,
- 75
- (7) D-3-(naphth-1-yl)alanyl,
 - (8) D-3-(naphth-2-yl)alanyl,
 - (9) D-(3-pyridyl)-alanyl,
 - (10) D-2-aminobutyryl,
 - (11) D-allo-isoleucyl,
 - (12) D-allo-threonyl,
- 80
- (13) D-allylglycyl,
 - (14) D-asparaginyl,
 - (15) D-aspartyl,
 - (16) D-benzothieryl,
 - (17) D-3-(4,4'-biphenyl)alanyl,
- 85
- (18) D-chlorophenylalanyl,
 - (19) D-3-(3-trifluoromethylphenyl)alanyl,
 - (20) D-3-(3-cyanophenyl)alanyl,
 - (21) D-3-(3,4-difluorophenyl)alanyl,
 - (22) D-citrullyl,
- 90
- (23) D-cyclohexylalanyl,
 - (24) D-cyclohexylglycyl,
 - (25) D-cystyl,
 - (26) D-cystyl(*S*-*t*-butyl),

95	(27)	D-glutaminyI,
	(28)	D-glutamyl,
	(29)	D-histidyl,
	(30)	D-homoisoleucyl,
	(31)	D-homophenylalanyl,
100	(32)	D-homoseryl,
	(33)	D-isoleucyl,
	(34)	D-leucyl,
	(35)	D-lysyl(N-epsilon-nicotinyl),
	(36)	D-lysyl,
105	(37)	D-methionyl,
	(38)	D-neopentylglycyl,
	(39)	D-norleucyl,
	(40)	D-norvalyl,
	(41)	D-ornithyl,
110	(42)	D-penicillaminyI,
	(43)	D-penicillaminyI(acetamidomethyl),
	(44)	D-penicillaminyI(S-benzyl),
	(45)	D-phenylalanyl,
	(46)	D-3-(4-aminophenyl)alanyl,
115	(47)	D-3-(4-methylphenyl)alanyl,
	(48)	D-3-(4-nitrophenyl)alanyl,
	(49)	D-3-(3,4-dimethoxyphenyl)alanyl,
	(50)	D-3-(3,4,5-trifluorophenyl)alanyl,
	(51)	D-prolyl,
120	(52)	D-seryl,
	(53)	D-seryl(O-benzyl),
	(54)	D-t-butylglycyl,
	(55)	D-thienylalanyl,
	(56)	D-threonyl,
125	(57)	D-threonyl(O-benzyl),
	(58)	D-tryptyl,

- (59) D-tyrosyl(*O*-benzyl),
- (60) D-tyrosyl(*O*-ethyl),
- (61) D-tyrosyl, and
- (62) D-valyl;

130 A₅ is an amino acyl residue of L or D configuration selected from:

- (1) alanyl,
- (2) (3-pyridyl)alanyl,
- (3) 3-(naphth-1-yl)alanyl,
- (4) 3-(naphth-2-yl)alanyl,
- (5) allo-threonyl,
- (6) allylglycyl,
- (7) glutaminyl,
- (8) glycyl,
- (9) histidyl,
- (10) homoseryl,
- (11) isoleucyl,
- (12) lysyl(N-epsilon-acetyl),
- (13) methionyl,
- (14) norvalyl,
- (15) octylglycyl,
- (16) ornithyl,
- (17) 3-(4-hydroxymethylphenyl)alanyl,
- (18) prolyl,
- (19) seryl,
- (20) threonyl,
- (21) tryptyl,
- (22) tyrosyl,
- (23) D-allo-threonyl,
- (24) D-homoseryl,
- (25) D-seryl,
- (26) D-threonyl,
- (27) penicillaminyl, and

(28) cystyl;

A₆ is an amino acyl residue of L or D configuration selected from:

- | | |
|-----|--------------------------------------|
| 160 | (1) alanyl, |
| | (2) 3-(naphth-1-yl)alanyl, |
| | (3) 3-(naphth-2-yl)alanyl, |
| | (4) (3-pyridyl)alanyl, |
| | (5) 2-aminobutyryl, |
| 165 | (6) allylglycyl, |
| | (7) arginyl, |
| | (8) asparaginyl, |
| | (9) aspartyl, |
| | (10) citrullyl, |
| 170 | (11) cyclohexylalanyl, |
| | (12) glutaminyl, |
| | (13) glutamyl, |
| | (14) glycyl, |
| | (15) histidyl, |
| 175 | (16) homoalanyl, |
| | (17) homoleucyl, |
| | (18) homoseryl, |
| | (19) isoleucyl, |
| | (20) leucyl, |
| 180 | (21) lysyl(N-epsilon-acetyl), |
| | (22) lysyl(N-epsilon-isopropyl), |
| | (23) methionyl(sulfone), |
| | (24) methionyl(sulfoxide), |
| | (25) methionyl, |
| 185 | (26) norleucyl, |
| | (27) norvalyl, |
| | (28) octylglycyl, |
| | (29) phenylalanyl, |
| | (30) 3-(4-carboxyamidophenyl)alanyl, |

- 190 (31) propargylglycyl,
 (32) seryl,
 (33) threonyl,
 (34) tryptyl,
 (35) tyrosyl,
- 195 (36) valyl,
 (37) D-3-(naphth-1-yl)alanyl,
 (38) D-3-(naphth-2-yl)alanyl,
 (39) D-glutaminyl,
 (40) D-homoseryl,
- 200 (41) D-leucyl,
 (42) D-norvalyl,
 (43) D-seryl,
 (44) penicillaminyl, and
 (45) cystyl;
- 205 A₇ is an amino acyl residue of L or D configuration selected from:
- (1) alanyl,
 (2) allylglycyl,
 (3) aspartyl,
 (4) citrullyl,
- 210 (5) cyclohexylglycyl,
 (6) glutamyl,
 (7) glycyl,
 (8) homoseryl,
 (9) isoleucyl,
- 215 (10) allo-isoleucyl
 (11) leucyl,
 (12) lysyl(N-epsilon-acetyl),
 (13) methionyl,
- 220 (14) 3-(naphth-1-yl)alanyl,
 (15) 3-(naphth-2-yl)alanyl,
 (16) norvalyl,

- 225 (17) phenylalanyl,
 (18) prolyl,
 (19) seryl,
 (20) *t*-butylglycyl,
 (21) tryptyl,
 (22) tyrosyl,
 (23) valyl,
 (24) D-allo-isoleucyl
 230 (25) D-isoleucyl,
 (26) penicillaminyl, and
 (27) cystyl;

A₈ is an amino acyl residue selected from:

- 235 (1) 2-amino-4-[(2-amino-pyrimidinyl)butanoyl,
 (2) alanyl(3-guanidino),
 (3) alanyl[3-pyrrolidiny(2-N-amidino)],
 (4) alanyl[4-piperidiny(N-amidino)],
 (5) arginyl,
 (6) arginyl(N^GN^Gdiethyl),
 240 (7) citrullyl,
 (8) 3-(cyclohexyl)alanyl(4-Nⁱ-isopropyl),
 (9) glycyl[4-piperidiny(N-amidino)],
 (10) histidyl,
 (11) homoarginyl,
 245 (12) lysyl,
 (13) lysyl(N-epsilon-isopropyl),
 (14) lysyl(N-epsilon-nicotinyl),
 (15) norarginyl,
 (16) ornithyl(N-delta-isopropyl),
 250 (17) ornithyl(N-delta-nicotinyl),
 (18) ornithyl[N-delta-(2-imidazoliny)],
 (19) [(4-amino(N-isopropyl)methyl)phenyl]alanyl,
 (20) 3-(4-guanidinophenyl)alanyl, and

(21) 3-(4-amino-N-isopropylphenyl)alanyl;

255

A₉ is an amino acyl residue of L or D configuration selected from:

- (1) 2-amino-butyryl,
- (2) 2-amino-isobutyryl,
- (3) homopropyl,
- (4) hydroxypropyl,
- (5) isoleucyl,
- (6) leucyl,
- (7) phenylalanyl,
- (8) prolyl,
- (9) seryl,

260

- (10) *t*-butylglycyl,
- (11) 1,2,3,4-tetrahydroisoquinoline-3-carbonyl,
- (12) threonyl,
- (13) valyl,
- (14) D-alanyl, and
- (15) D-prolyl; and

265

270


A₁₀ is a hydroxyl group or an amino acid amide is selected from:

- (1) azaglycyclamide,
- (2) D-alanylamide,
- (3) D-alanylethylamide,
- (4) glycylamide,
- (5) glycylethylamide,
- (6) sarcosylamide,
- (7) serylamide,
- (8) D-serylamide,

275

280

- (9) a group represented by the formula



- (10) a group represented by the formula -NH-R⁴ ;

wherein:

s is an integer selected from 0 to 8,

285 R² is selected from hydrogen, alkyl, and a 5- to 6-membered cycloalkyl ring;

R³ is selected from hydrogen, hydroxy, alkyl, phenyl, alkoxy, and a 5- to 6-membered ring optionally containing from one to two heteroatoms selected from oxygen, nitrogen, and sulfur, provided that s is not zero when R³ is hydroxy or alkoxy; and

290 R⁴ is selected from hydrogen, hydroxy, and a 5- to 6-membered cycloalkyl ring.

2. A compound according to Claim 1, wherein A₁ is sarcosyl, A₂ is glycyl, A₃ is valyl, A₇ is isoleucyl, A₈ is arginyl, A₉ is prolyl, and A₀, A₄, A₅, A₆, and A₁₀ are as defined in Claim 1.

3. A compound according to Claim 2, wherein A₄ is an amino acyl residue having a D configuration selected from:

- 5 (1) D-alanyl,
(2) D-3-(naphth-1-yl)alanyl,
(3) D-3-(naphth-2-yl)alanyl,
(4) D-(3-pyridyl)-alanyl,
(5) D-2-aminobutyryl,
(6) D-allo-isoleucyl,
10 (7) D-allo-threonyl,
(8) D-allylglycyl,
(9) D-asparaginy, l,
(10) D-aspartyl,
(11) D-chlorophenylalanyl,
15 (12) D-3-(3-trifluoromethylphenyl)alanyl,
(13) D-3-(3-cyanophenyl)alanyl,
(14) D-3-(3,4-difluorophenyl)alanyl,
(15) D-cyclohexylalanyl,

20	(16) D-cyclohexylglycyl,
	(17) D-cystyl,
	(18) D-glutaminyI,
	(19) D-glutamyl,
	(20) D-histidyl,
	(21) D-homoisoleucyl,
25	(22) D-homophenylalanyl,
	(23) D-homoseryl,
	(24) D-isoleucyl,
	(25) D-leucyl,
	(26) D-lysyl(N-epsilon-nicotinyl),
30	(27) D-methionyl,
	(28) D-neopentylglycyl,
	(29) D-norleucyl,
	(30) D-norvalyl,
	(31) D-penicillaminyI,
35	(32) D-penicillaminyI(acetamidomethyl),
	(33) D-penicillaminyI(S-benzyl),
	(34) D-phenylalanyl,
	(35) D-3-(4-aminophenyl)alanyl,
	(36) D-3-(4-methylphenyl)alanyl,
40	(37) D-3-(4-nitrophenyl)alanyl,
	(38) D-3-(3,4-dimethoxyphenyl)alanyl,
	(39) D-3-(3,4,5-trifluorophenyl)alanyl,
	(40) D-prolyl,
	(41) D-seryl,
45	(42) D-seryl(<i>O</i> -benzyl),
	(43) D- <i>t</i> -butylglycyl,
	(44) D-thienylalanyl,
	(45) D-threonyl,
	(46) D-threonyl(<i>O</i> -benzyl),
50	(47) D-tyrosyl(<i>O</i> -ethyl),

(48) D-tyrosyl, and

(49) D-valyl.

4. A compound according to Claim 3, wherein A₄ is an amino acyl residue having a D configuration selected from:

(1) D-allo-isoleucyl,

(2) D-allylglycyl,

(3) D-3-(3-cyanophenyl)alanyl,

(4) D-cystyl,

(5) D-isoleucyl,

(6) D-leucyl,

(7) D-penicillaminy,

(8) D-phenylalanyl,

(9) D-3-(3,4,5-trifluorophenyl)alanyl, and

(10) D-3-(4-aminophenyl)alanyl.

5. A compound according to Claim 2, wherein A₅ is selected from:

(1) glycyl,

(2) octylglycyl,

(3) penicillaminy,

(4) seryl,

(5) threonyl, and

(6) tyrosyl.

6. A compound according to Claim 2, wherein A₆ is selected from:

(1) glutaminy,

(2) leucyl,

(3) norvalyl, and

(4) seryl.

7. A compound according to Claim 3, wherein A_0 is selected from:

- (1) acetyl,
- (2) butyryl,
- (3) caproyl,
- (4) (4-N-acetylamino)butyryl,
- (5) N-acetyl-beta-alanyl,
- (6) (6-N-acetylamino)caproyl,
- (7) chloronicotinyl,
- (8) cyclohexylacetyl,
- (9) furoyl,
- (10) gamma-aminobutyryl,
- (11) 2-methoxyacetyl,
- (12) methylnicotinyl,
- (13) nicotinyl,
- (14) (8-N-acetylamino)-3,6-dioxo-octanoyl,
- (15) phenylacetyl,
- (16) propionyl,
- (17) shikimyl,
- (18) succinyl, and
- (19) tetrahydrofuroyl.

8. A compound according to Claim 3, wherein A_{10} is selected from:

- (1) D-alanylamine,
- (2) azaglycylamine,
- (3) serylamine,
- (4) ethylamine,
- (5) hydroxylamine,
- (6) isopropylamine,
- (7) propylamine,
- (8) 2-(cyclohexyl)ethylamine,
- (9) 2-(1-pyrrolidine)ethylamine,
- (10) 1-(cyclohexyl)ethylamine,

- 15 (11) 2-(methoxy)ethylamide,
 (12) 2-(hydroxy)ethylamide,
 (13) 2-(2-pyridine)ethylamide,
 (14) (2-pyridine)methylamide,
 (15) 2-(3-pyridine)ethylamide,
 (16) 2-(2-(1-methyl)pyrrolidine)ethylamide,
 (17) 2-(N-morpholine)ethylamide, and
 20 (18) cyclopropylmethylamide.

9. A compound according to Claim 1, wherein A₄ is an amino acyl residue having a D configuration selected from:

- 5 (1) D-allo-isoleucyl,
 (2) D-allylglycyl,
 (3) D-3-(3-cyanophenyl)alanyl,
 (4) D-cystyl,
 (5) D-isoleucyl,
 (6) D-leucyl,
 10 (7) D-penicillaminyl,
 (8) D-phenylalanyl,
 (9) D-3-(3,4,5-trifluorophenyl)alanyl, and
 (10) D-3-(4-aminophenyl)alanyl;

A₅ is an amino acyl residue selected from:

- 15 (1) octylglycyl,
 (2) glycyl,
 (3) penicillaminyl,
 (4) seryl,
 (5) threonyl, and
 20 (6) tyrosyl; and

A₆ is an amino acyl residue selected from:

- (1) glutaminyl,
 (2) leucyl,
 (3) norvalyl, and

25

(4) seryl.

10. A compound according to Claim 9, wherein A_0 is selected from:

(1) acetyl,

(2) butyryl,

5

(3) caproyl,

(4) (4-N-acetylamino)butyryl,

(5) N-acetyl-beta-alanyl,

(6) (6-N-acetylamino)caproyl,

(7) chloronicotinyl,

10

(8) cyclohexylacetyl,

(9) furoyl,

(10) gamma-aminobutyryl,

(11) 2-methoxyacetyl,

(12) methylnicotinyl,

15

(13) nicotinyl,

(14) (8-N-acetylamino)-3,6-dioxo-octanoyl,

(15) phenylacetyl,

(16) propionyl,

(17) shikimyl,

20

(18) succinyl, and

(19) tetrahydrofuroyl.

11. A compound according to Claim 9, wherein A_{10} is selected from:

25

(1) D-alanylamide,

(2) azaglycylamide,

(3) serylamine

(4) ethylamide,

(5) hydroxylamide,

30

(6) isopropylamide,

(7) propylamide,

- 5 (8) 2-(cyclohexyl)ethylamide,
 (9) 2-(1-pyrrolidine)ethylamide,
 (10) 1-(cyclohexyl)ethylamide,
 (11) 2-(methoxy)ethylamide,
 (12) 2-(hydroxy)ethylamide,
 (13) 2-(2-pyridine)ethylamide,
 (14) (2-pyridine)methylamide,
 (15) 2-(3-pyridine)ethylamide,
 (16) 2-(2-(1-methyl)pyrrolidine)ethylamide,
 10 (17) 2-(N-morpholine)ethylamide, and
 (18) cyclopropylmethylamide.

12. A compound, or a pharmaceutically acceptable salt, ester, solvate or prodrug thereof, selected from:

- 15 (1) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (2) pyroGlu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (3) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₃,
 (4) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (5) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂-(1-pyrrolidine),
 20 (6) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethylpiperidine,
 (7) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHmethylcyclopropyl,
 (8) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH(ethyl-1-(R)-cyclohexyl),
 (9) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂,
 (10) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 25 (11) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂cyclohexyl,
 (12) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (13) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (14) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (15) N-Ac-Sar-Gly-Val-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 (16) N-Ac-Sar-Gly-Val-Gly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (17) N-Ac-Sar-Gly-Val-D-Val-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (18) N-Ac-Sar-Gly-Val-D-Ala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (19) N-Ac-Sar-Gly-Val-D-Met-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- (20) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (21) N-Ac-Sar-Gly-Val-D-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (22) N-Ac-Sar-Gly-Val-D-Tyr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (23) N-Ac-Sar-Gly-Val-D-4,4'-Biphenylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (24) N-Ac-Sar-Gly-Val-D-Cha-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (25) N-Ac-Sar-Gly-Val-D-Chg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (26) N-Ac-Sar-Gly-Val-D-4-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (27) N-Ac-Sar-Gly-Val-D-Hphe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (28) N-Ac-Sar-Gly-Val-Dehydroleu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 (29) N-Ac-Sar-Gly-Val-D-3-CF₃Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (30) N-Ac-Sar-Gly-Val-D-pentaFphe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (31) N-Ac-Sar-Gly-Val-D-3,4-diClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (32) N-Ac-Sar-Gly-Val-D-3-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (33) N-Ac-Sar-Gly-Val-D-2-Thienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 (34) N-Ac-Sar-Gly-Val-D-3-CNphe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (35) N-Ac-Sar-Gly-Val-D-Ile-Thr-DNva-Ile-Arg-ProNHCH₂CH₃,
 (36) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (37) N-Ac-Sar-Gly-Val-D-Ile-Thr-Cha-Ile-Arg-ProNHCH₂CH₃,
 (38) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gly-Ile-Arg-ProNHCH₂CH₃,
 20 (39) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂CH₃,
 (40) N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-ProNHCH₂CH₃,
 (41) N-Ac-Sar-Gly-Val-D-Ile-Thr-Abu-Ile-Arg-ProNHCH₂CH₃,
 (42) N-Ac-Sar-Gly-Val-D-Ile-Thr-Allylgly-Ile-Arg-ProNHCH₂CH₃,
 (43) N-Ac-Sar-Gly-Val-D-Ile-Thr-Octylgly-Ile-Arg-ProNHCH₂CH₃,
 25 (44) N-Ac-Sar-Gly-Val-D-Ile-Thr-Met-Ile-Arg-ProNHCH₂CH₃,
 (45) N-Cyclohexylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (46) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (47) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (48) N-Nicotinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 (49) N-Propionyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (50) N-(MeO)acetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (51) N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (52) N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- (53) N-Butyryl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (54) N[2-THFcarbonyl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (55) N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (56) N[6-N-acetyl-(CH₂)₅C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (57) N-Hexanoyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (58) N-[4-N-Acetylaminobutyryl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 (59) H-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (60) N-Ac-Sar-Gly-Asn-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (61) N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (62) N-Ac-Pro-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 (63) N-Ac-Gly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (64) N-Ac-Ala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (65) N-Ac-NEtGly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (66) N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (67) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 20 (68) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (69) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-ProNHCH₂CH₃,
 (70) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-AbuNHCH₂CH₃,
 (71) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Phe-NHCH₂CH₃,
 (72) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Tic-NHCH₂CH₃,
 25 (73) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Hyp-NHCH₂CH₃,
 (74) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Aib-NHCH₂CH₃,
 (75) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-D-Ala-NHCH₂CH₃,
 (76) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pip-NHCH₂CH₃,
 (77) N-Ac-Sar-Gly-Val-D-Tyr(Et)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 (78) N-Ac-Sar-Gly-Val-D-Cys(tBu)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (79) N-Ac-Sar-Gly-Val-D-Cys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (80) N-Ac-Sar-Gly-Val-D-Tyr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (81) N-Ac-Sar-Gly-Val-D-Ser(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (82) N-Ac-Sar-Gly-Val-D-INal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- (83) N-Ac-Sar-Gly-Val-D-tButylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (84) N-Ac-Sar-Gly-Val-D-Orn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (85) N-Ac-Sar-Gly-Val-D-Thr(Bzl)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (86) N-Ac-Sar-Gly-Val-D-2Nal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (87) N-Ac-Sar-Gly-Val-D-Phe(4-Me)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (88) N-Ac-Sar-Gly-Val-D-Phe(3,4-diMeO)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (89) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (90) N-Ac-Sar-Gly-Val-D-Phe(4-NO₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (91) N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 (92) N-Ac-Sar-Gly-Val-D-Pen(Acm)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (93) N-Ac-Sar-Gly-Val-D-Abu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (94) N-Ac-Sar-Gly-Val-D-Phe(4-NH₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (95) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ala-Arg-ProNHCH₂CH₃,
 (96) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Met-Arg-ProNHCH₂CH₃,
 15 (97) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Phe-Arg-ProNHCH₂CH₃,
 (98) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Tyr-Arg-ProNHCH₂CH₃,
 (99) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Nva-Arg-ProNHCH₂CH₃,
 (100) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Asp-Arg-ProNHCH₂CH₃,
 (101) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Gly-Arg-ProNHCH₂CH₃,
 20 (102) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Lys(Ac)-Arg-ProNHCH₂CH₃,
 (103) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Leu-Arg-ProNHCH₂CH₃,
 (104) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-2Nal-Arg-ProNHCH₂CH₃,
 (105) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-1Nal-Arg-ProNHCH₂CH₃,
 (106) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Allylgly-Arg-ProNHCH₂CH₃,
 25 (107) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Cit-Arg-ProNHCH₂CH₃,
 (108) N-Ac-Sar-Gly-Val-D-Leu-Ala-Nva-Ile-Arg-ProNHCH₂CH₃,
 (109) N-Ac-Sar-Gly-Val-D-Leu-Pro-Nva-Ile-Arg-ProNHCH₂CH₃,
 (110) N-Ac-Sar-Gly-Val-D-Leu-Trp-Nva-Ile-Arg-ProNHCH₂CH₃,
 (111) N-Ac-Sar-Gly-Val-D-Leu-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 (112) N-Ac-Sar-Gly-Val-D-Leu-Nva-Nva-Ile-Arg-ProNHCH₂CH₃,
 (113) N-Ac-Sar-Gly-Val-D-Leu-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (114) N-Ac-Sar-Gly-Val-D-Leu-Lys(Ac)-Nva-Ile-Arg-ProNHCH₂CH₃,
 (115) N-Ac-Sar-Gly-Val-D-Leu-2Nal-Nva-Ile-Arg-ProNHCH₂CH₃,

- (116) N-Ac-Sar-Gly-Val-D-Leu-1Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
 (117) N-Ac-Sar-Gly-Val-D-Leu-Octylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (118) N-Ac-Sar-Gly-Val-D-Leu-Gln-Nva-Ile-Arg-ProNHCH₂CH₃,
 (119) N-Ac-Sar-Gly-Val-D-Leu-Met-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (120) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (121) N-Ac-Sar-Gly-Val-D-Leu-Allylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (122) N-Ac-Sar-Gly-Val-D-Leu-Ile-Nva-Ile-Arg-ProNHCH₂CH₃,
 (123) N-Ac-Sar-Gly-Val-D-Leu-D-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (124) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ile-Ile-Arg-ProNHCH₂CH₃,
 10 (125) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nle-Ile-Arg-ProNHCH₂CH₃,
 (126) N-Ac-Sar-Gly-Val-D-Ile-Thr-Cit-Ile-Arg-ProNHCH₂CH₃,
 (127) N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O₂)-Ile-Arg-ProNHCH₂CH₃,
 (128) N-Ac-Sar-Gly-Val-D-Ile-Thr-Arg-Ile-Arg-ProNHCH₂CH₃,
 (129) N-Ac-Sar-Gly-Val-D-Ile-Thr-Tyr-Ile-Arg-ProNHCH₂CH₃,
 15 (130) N-Ac-Sar-Gly-Val-D-Ile-Thr-Glu-Ile-Arg-ProNHCH₂CH₃,
 (131) N-Ac-Sar-Gly-Val-D-Ile-Thr-Lys(Ac)-Ile-Arg-ProNHCH₂CH₃,
 (132) N-Ac-Sar-Gly-Val-D-Ile-Thr-Propargylgly-Ile-Arg-ProNHCH₂CH₃,
 (133) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (134) N-Ac-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 20 (135) N-Ac-Bala-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (136) N-Phenylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (137) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂,
 (138) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Sar-NHCH₂CH₃,
 (139) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SerNH₂,
 25 (140) N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (141) N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (142) N-Ac-Sar-Leu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (143) N-Ac-Sar-Phe-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (144) N-Ac-Sar-Glu-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 (145) N-Ac-Sar-Pro-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (146) N-Ac-Sar-Asn-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (147) N-Ac-Sar-Asp-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (148) N-Ac-Asn-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- (149) N-Ac-Gln-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (150) N-Ac-Ser-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (151) N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (152) N-Ac-Glu-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (153) N-Ac-Gaba-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (154) N-Ac-Bala-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (155) N-Ac-Gln-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (156) N-Ac-Sar-Gly-Gly-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (157) N-Ac-Sar-Gly-Glu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 (158) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (159) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (160) N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (161) N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (162) N-Ac-Sar-Gly-Val-D-Leu-Thr-Asp-Ile-Arg-ProNHCH₂CH₃,
 15 (163) N-Ac-Sar-Gly-Val-D-Ile-Thr-Asp-Ile-Arg-ProNHCH₂CH₃,
 (164) N-Ac-Sar-Gly-Val-D-Ile-Thr-Asn-Ile-Arg-ProNHCH₂CH₃,
 (165) N-Ac-Sar-Gly-Val-D-Ile-Thr-Met(O)-Ile-Arg-ProNHCH₂CH₃,
 (166) N-Ac-Sar-Gly-Val-D-Leu-Thr-Asn-Ile-Arg-ProNHCH₂CH₃,
 (167) N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (168) N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (169) N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (170) N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (171) N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (172) N-Ac-Sar-Gly-Val-D-Cit-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 (173) N-Ac-Sar-Gly-Val-D-Hcit-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (174) N-Ac-Sar-Gly-Val-D-Hle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (175) N-Ac-Sar-Gly-Val-D-Neopentylgly-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (176) N-Ac-Sar-Gly-Val-D-Ile-Thr-Phe(4-CONH₂)-Ile-Arg-ProNHCH₂CH₃,
 (177) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-His-ProNHCH₂CH₃,
 30 (178) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Isp)-ProNHCH₂CH₃,
 (179) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys(Nic)-ProNHCH₂CH₃,
 (180) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Om(Nic)-ProNHCH₂CH₃,
 (181) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Om(Isp)-ProNHCH₂CH₃,

- (182) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-Nlsp)-ProNHCH₂CH₃,
 (183) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cha(4-Nlsp)-ProNHCH₂CH₃,
 (184) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Harg-ProNHCH₂CH₃,
 (185) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Norarg-ProNHCH₂CH₃,
 5 (186) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Cit-ProNHCH₂CH₃,
 (187) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Lys-ProNHCH₂CH₃,
 (188) N-Ac-Sar-Gly-Val-D-Ile-Phe(4-CH₂OH)-Nva-Ile-Arg-ProNHCH₂CH₃,
 (189) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-guanidino)-ProNHCH₂CH₃,
 10 (190) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Aminopyrimidinylbutanoyl-Pro-
 NHCH₂CH₃,
 (191) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Phe(4-CH₂NHlsp)-ProNHCH₂CH₃,
 (192) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Gly[4-Pip(N-amidino)]-Pro-
 NHCH₂CH₃,
 15 (193) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala[4-Pip(N-amidino)]-Pro-
 NHCH₂CH₃,
 (194) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-guanidino)-ProNHCH₂CH₃,
 (195) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Ala(3-pyrrolidinylamidino)-Pro-
 NHCH₂CH₃,
 20 (196) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Orn(2-imidazo)-ProNHCH₂CH₃,
 (197) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (198) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (199) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (200) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 25 (201) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (202) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (203) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (204) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (205) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 30 (206) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (207) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (208) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (209) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (210) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SarNH₂,
 (211) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-SarNH₂,

- (212) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-SarNH₂,
 (213) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-SarNH₂,
 (214) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-Pro-D-AlaNH₂,
 (215) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂(CH₃)₂,
 5 (216) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 (217) N-Ac-Sar-Gly-Val-D-Ile-Thr-Orn(Ac)-Ile-Arg-ProNHCH₂CH₃,
 (218) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂,
 (219) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂,
 (220) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-AzaglyNH₂,
 10 (221) N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-
 NHCH₂CH₃,
 (222) N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (223) N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-
 NHCH₂CH₃,
 15 (224) N-(2-THFcarbonyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (225) N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-
 AlaNH₂,
 (226) N-(2-THFcarbonyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-
 NHCH₂(CH₃)₂,
 20 (227) N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (228) N-(6-Ac-Aca)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (229) N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (230) N-(6-Ac-Aca)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (231) N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 25 (232) N-(6-Ac-Aca)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (233) N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (234) N-(4-Ac-Gaba)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (235) N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (236) N-(4-Ac-Gaba)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 30 (237) N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (238) N-(4-Ac-Gaba)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-
 NHCH₂(CH₃)₂,
 (239) N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (240) N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 35 (241) N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

- (242) N-(2-Furoyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (243) N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (244) N-(2-Furoyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (245) N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (246) N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (247) N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (248) N-(Shikimyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (249) N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (250) N-(Shikimyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 10 (251) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-
 NHCH₂CH₃,
 (252) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (253) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-
 NHCH₂CH₃,
 15 (254) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-D-AlaNH₂,
 (255) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-D-
 AlaNH₂,
 (256) N-(2-Me-Nicotinyl)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-
 NHCH₂(CH₃)₂,
 20 (257) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 (258) N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂,
 (259) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (260) N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 (261) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 25 (262) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂(CH₃)₂,
 (263) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (264) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (265) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Leu-Ile-Arg-Pro-D-AlaNH₂,
 (266) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-Pro-AzaglyNH₂,
 30 (267) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHethyl-(1-pyrrolidine),
 (268) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNH(ethyl-1-cyclohexyl),
 (269) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHethyl-(1-pyrrolidine),
 (270) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-cyclohexyl),
 35 (271) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNH(ethyl-1-
 cyclohexyl),

- (272) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (273) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (274) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (275) N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₂OCH₃,
 5 (276) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (277) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (278) N-Succinyl-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (279) N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 (280) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₂OCH₃,
 10 (281) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Allygly-Ile-Arg-ProNHCH₂CH₃,
 (282) N-Ac-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-ProNHCH₂(CH₃)₂,
 (283) N-Ac-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 (284) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 (285) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Allygly-Ile-Arg-Pro-D-AlaNH₂,
 15 (286) N-Ac-Sar-Gly-Val-D-Ile-Ser-Allygly-Ile-Arg-Pro-ProNHCH₂CH₃,
 (287) N-Ac-Sar-Gly-Val-D-Leu-Ser-Allygly-Ile-Arg-Pro-ProNHCH₂CH₃,
 (288) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SarNH₂,
 (289) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂,
 (290) N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (291) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (292) N-Ac-Sar-Gly-Val-D-Leu-Hser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (293) N-Ac-Sar-Gly-Gln-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (294) N-Ac-Sar-Gly-Nva-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (295) N-Ac-Sar-Gly-Ile-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 (296) N-Ac-Sar-Gly-Phe-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (297) N-Ac-Sar-Gly-Leu-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (298) N-Ac-Sar-Gly-Ser-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (299) N-Ac-Thr-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (300) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ala-Ile-Arg-ProNHCH₂CH₃,
 30 (301) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-ProNHCH₂(CH₃)₂,
 (302) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,
 (303) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,
 (304) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Ala-Ile-Arg-Pro-D-AlaNH₂,

- (305) N-Ac-Sar-Gly-Val-D-Ile-Ser-Ala-Ile-Arg-ProNHCH₂CH₃,
 (306) N-Ac-Sar-Gly-Val-D-Leu-Ser-Ala-Ile-Arg-ProNHCH₂CH₃,
 (307) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Val-Ile-Arg-ProNHCH₂CH₃,
 (308) N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-ProNHCH₂(CH₃)₂,
 5 (309) N-Ac-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,
 (310) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,
 (311) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Val-Ile-Arg-Pro-D-AlaNH₂,
 (312) N-Ac-Sar-Gly-Val-D-Ile-Ser-Val-Ile-Arg-ProNHCH₂CH₃,
 (313) N-Ac-Sar-Gly-Val-D-Leu-Ser-Val-Ile-Arg-ProNHCH₂CH₃,
 10 (314) N-Ac-Sar-Gly-Val-D-alloIle-Thr-D-Nva-Ile-Arg-ProNHCH₂CH₃,
 (315) N-Ac-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (316) N-Ac-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (317) N-Ac-Sar-Gly-Val-D-alloIle-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (318) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-D-Nva-Ile-Arg-Pro-D-AlaNH₂,
 15 (319) N-Ac-Sar-Gly-Val-D-Ile-Ser-D-Nva-Ile-Arg-ProNHCH₂CH₃,
 (320) N-Ac-Sar-Gly-Val-D-Leu-Ser-D-Nva-Ile-Arg-ProNHCH₂CH₃,
 (321) N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (322) N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (323) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 20 (324) N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (325) N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (326) N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (327) N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (328) N-Succinyl-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 25 (329) N-Ac-Sar-Gly-Val-D-Ile-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (330) N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (331) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (332) N-Ac-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (333) N-Ac-Sar-Gly-Val-D-Leu-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 30 (334) N-Ac-Sar-Gly-Val-D-Ile-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 (335) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (336) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (337) N-Succinyl-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,

- (338) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (339) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (340) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 (341) N-Ac-Sar-Gly-Val-D-alloIle-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 5 (342) N-Ac-Sar-Gly-Val-D-Ile-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (343) N-Ac-Sar-Gly-Val-D-alloIle-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (344) N-Ac-Sar-Gly-Val-D-Leu-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 (345) N-Ac-Sar-Gly-Val-D-Ile-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 (346) N-Ac-Sar-Gly-Val-D-alloIle-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 10 (347) N-Ac-Sar-Gly-Val-D-Ile-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (348) N-Ac-Sar-Gly-Val-D-alloIle-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (349) N-Ac-Sar-Gly-Val-D-Leu-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (350) N-Ac-Sar-Gly-Val-D-Ile-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (351) N-Ac-Sar-Gly-Val-D-alloIle-Tyr-Gln-Ile-Arg-ProNHCH₂CH₃,
 15 (352) N-Ac-Sar-Gly-Val-D-Ser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (353) N-Ac-Sar-Gly-Val-D-Thr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (354) N-Ac-Sar-Gly-Val-D-Gln-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (355) N-Ac-Sar-Gly-Val-D-Asn-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (356) N-Ac-Sar-Gly-Val-D-Arg-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (357) N-Ac-Sar-Gly-Val-D-3-Pal-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (358) N-Ac-Sar-Gly-Val-D-Glu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (359) N-Ac-Sar-Gly-Val-D-Asp-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (360) N-Ac-Sar-Gly-Val-D-His-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (361) N-Ac-Sar-Gly-Val-D-Hser-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 (362) N-Ac-Sar-Gly-Val-D-alloThr-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (363) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-D-Ile-Arg-ProNHCH₂CH₃,
 (364) N-Ac-Sar-Gly-Val-D-Ser-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (365) N-Ac-Sar-Gly-Val-D-Thr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (366) N-Ac-Sar-Gly-Val-D-alloThr-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 30 (367) N-Ac-Sar-Gly-Val-D-Ser-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (368) N-Ac-Sar-Gly-Val-D-Thr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (369) N-Ac-Sar-Gly-Val-D-alloThr-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (370) N-Ac-Sar-Gly-Val-D-alloThr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,

- (371) N-Ac-Sar-Gly-Val-D-Thr-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (372) N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (373) N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (374) N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 5 (375) N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (376) N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (377) N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (378) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (379) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 10 (380) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (381) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (382) N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 15 (383) N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (384) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 (385) N-Ac-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 (386) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 (387) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 20 (388) N-Ac-Sar-Gly-Val-D-Leu-Ser-Ser-Ile-Arg-ProNHethyl-1-(R)-cyclohexyl,
 (389) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHethyl-1-(S)-cyclohexyl,
 (390) N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (391) N-Ac-Sar-Gly-Val-D-Pen-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (392) N-Ac-Sar-Gly-Val-D-Pen-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 25 (393) N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (394) N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (395) N-Ac-Sar-Gly-Val-D-Pen-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (396) N-Ac-Sar-Gly-Val-D-Pen-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (397) N-Ac-Sar-Gly-Val-D-Pen-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 30 (398) N-Ac-Sar-Gly-Val-D-Pen-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (399) N-Ac-Sar-Gly-Val-D-Pen-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 (400) N-Ac-Sar-Gly-Val-D-Pen-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (401) N-Ac-Sar-Gly-Val-D-Pen-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,

- (402) N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (403) N-Succinyl-Sar-Gly-Val-D-Pen-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 (404) N-Succinyl-Sar-Gly-Val-D-Pen-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (405) N-Ac-Sar-Gly-Val-D-Cys-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (406) N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (407) N-Ac-Sar-Gly-Val-D-Cys-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (408) N-Ac-Sar-Gly-Val-D-Cys-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (409) N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (410) N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 (411) N-Ac-Sar-Gly-Val-D-Cys-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (412) N-Ac-Sar-Gly-Val-D-Cys-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (413) N-Ac-Sar-Gly-Val-D-Cys-Gly-Gln-Ile-Arg-ProNHCH₂CH₃,
 (414) N-Ac-Sar-Gly-Val-D-Cys-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (415) N-Ac-Sar-Gly-Val-D-Cys-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 15 (416) N-Ac-Sar-Gly-Val-D-Cys-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (417) N-Ac-Sar-Gly-Val-D-Cys-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 (418) N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (419) N-Succinyl-Sar-Gly-Val-D-Cys-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 (420) N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (421) N-Ac-Sar-Gly-Cys-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (422) N-Ac-Sar-Gly-Pen-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (423) N-Ac-Sar-Gly-Pen-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (424) N-Ac-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (425) N-Ac-Sar-Gly-Pen-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 (426) N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (427) N-Ac-Sar-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (428) N-Succinyl-Gly-Pen-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (429) N-Succinyl-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (430) N-Succinyl-Sar-Gly-Pen-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 30 (431) N-Ac-Sar-Gly-Val-D-Leu-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,
 (432) N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,
 (433) N-Ac-Sar-Gly-Val-D-alloIle-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,

- (434) N-Ac-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂CH₃,
 (435) N-Ac-Sar-Gly-Val-D-Ile-Pen-Ser-Ile-Arg-ProNHCH₂CH₃,
 (436) N-Ac-Sar-Gly-Val-D-Ile-Pen-Leu-Ile-Arg-ProNHCH₂CH₃,
 (437) N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 5 (438) N-Ac-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (439) N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Nva-Ile-Arg-ProNHCH₂CH₃,
 (440) N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂CH₃,
 (441) N-Succinyl-Sar-Gly-Val-D-Ile-Pen-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (442) N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 10 (443) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 (444) N-Ac-Sar-Gly-Val-D-Leu-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 (445) N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-Pro-D-AlaNH₂,
 (446) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂CH₃,
 (447) N-Ac-Sar-Gly-Val-D-Ile-Thr-Pen-Ile-Arg-ProNHCH₂(CH₃)₂,
 15 (448) N-Ac-Sar-Gly-Val-D-Leu-Ser-Pen-Ile-Arg-ProNHCH₂CH₃,
 (449) N-Ac-Sar-Gly-Val-D-Leu-Gly-Pen-Ile-Arg-ProNHCH₂CH₃,
 (450) N-Succinyl-Sar-Gly-Val-D-Leu-Ser-Pen-Ile-Arg-ProNHCH₂CH₃,
 (451) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (452) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (453) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (454) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Leu-Ile-Arg-ProNHCH₂CH₃,
 (455) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (456) N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 25 (457) N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (458) N-Succinyl-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Gln-Ile-Arg-ProNH-CH₂(CH₃)₂,
 (459) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 30 (460) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Ser-Ser-Ile-Arg-ProNHCH₂CH₃,
 (461) N-Ac-Sar-Ala-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (462) N-Ac-Sar-Ala-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (463) N-Ac-Sar-Ala-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,

- (464) N-Ac-Sar-Ala-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (465) N-Ac-Sar-Ala-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (466) N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (467) N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-ProNHCH₂CH₃,
 5 (468) N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (469) N-Succinyl-Sar-Ala-Val-D-Ile-Thr-Gln-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (470) N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (471) N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (472) N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 10 (473) N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-DAlaNH₂,
 (474) N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-Pro-DAlaNH₂,
 (475) N-(3-Ac-Bala)-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (476) N-(3-Ac-Bala)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (477) N-(3-Ac-Bala)-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 15 (478) N-(3-Ac-Bala)-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (479) N-(3-Ac-Bala)-Sar-Gly-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (480) N-(3-Ac-Bala)-Sar-Ala-Val-D-alloIle-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (481) N-(3-Ac-Bala)-Sar-Ala-Val-D-Ile-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (482) N-(3-Ac-Bala)-Sar-Ala-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (483) N-(3-Ac-Bala)-Sar-Ala-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂CH₃,
 (484) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 (485) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-Pro-OH,
 (486) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Ile-Arg-Pro-OH,
 (487) N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-Pro-OH,
 25 (488) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-Pro-OH,
 (489) N-Ac-Sar-Gly-Val-D-Ile-Thr-Gln-Ile-Arg-Pro-OH,
 (490) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-Pro-OH,
 (491) N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 (492) N-Ac-Sar-Gly-Val-D-Ile-Ser-Gln-Ile-Arg-Pro-OH,
 30 (493) N-Succinyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 (494) N-Succinyl-Sar-Gly-Val-D-Leu-Thr-Gln-Ile-Arg-Pro-OH,
 (495) N-Ac-Sar-Gly-Asp-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- (496) N-Ac-Sar-Gly-Ala-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (497) N-Ac-Sar-Gly-Cha-D-Leu-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (498) N-Ac-Sar-Gly-Met-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (499) N-Ac-Cit-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃, N-Ac-Sar-Gly-Val-D-Ile-Thr-Hser-Ile-Arg-ProNHCH₂CH₃,
 (501) N-Ac-Sar-Gly-Val-DalloIle-His-Nva-Ile-Arg-ProNHCH₂CH₃,
 (502) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-Butyl,
 (503) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Butyl,
 (504) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-iso-Amyl,
 (505) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-n-hexyl,
 (506) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3,3-dimethyl)butyl,
 (507) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-ethoxy)ethyl,
 (508) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(2-isopropoxy)ethyl,
 (509) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(3-methoxy)propyl,
 (510) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-(cyclopentyl)methyl,
 and
 (511) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-cyclohexyl.
13. A compound according to Claim 12, selected from:
- (1) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (2) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₂-(1-pyrrolidine),
 (3) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH(ethyl-1-(R)-cyclohexyl),
 (4) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH₂,
 (5) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (6) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (7) N-Ac-Sar-Gly-Val-D-Val-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (8) N-Ac-Sar-Gly-Val-D-Nle-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (9) N-Ac-Sar-Gly-Val-D-Phe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (10) N-Ac-Sar-Gly-Val-D-Cha-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (11) N-Ac-Sar-Gly-Val-D-3,4-diClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (12) N-Ac-Sar-Gly-Val-D-3-ClPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (13) N-Ac-Sar-Gly-Val-D-2-Thienylala-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- 5 (14) N-Ac-Sar-Gly-Val-D-3-CNPhe-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (15) N-Ac-Sar-Gly-Val-D-Ile-Thr-Cha-Ile-Arg-ProNHCH₂CH₃,
 (16) N[2-THF-C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (17) N[6-N-acetyl-(CH₂)₅C(O)]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (18) N-Hexanoyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (19) N-[4-N-Acetylaminobutryl]-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 10 (20) N-[CH₃C(O)NH-(CH₂)₂-O-(CH₂)₂-O-CH₂-C(O)]-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (21) N-Ac-Pro-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (22) N-Ac-NEtGly-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (23) N-Ac-Sar-Gly-Val-D-Ile-Thr-Leu-Ile-Arg-ProNHCH₂CH₃,
 (24) N-Ac-Sar-Gly-Val-D-Ile-Thr-Ser-Ile-Arg-ProNHCH₂CH₃,
 15 (25) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
 (26) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Lys(Ac)-Arg-ProNHCH₂CH₃,
 (27) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Leu-Arg-ProNHCH₂CH₃,
 (28) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-1Nal-Arg-ProNHCH₂CH₃,
 (29) N-Ac-Sar-Gly-Val-D-Leu-Thr-Nva-Allylgly-Arg-ProNHCH₂CH₃,
 20 (30) N-Ac-Sar-Gly-Val-D-Leu-Ala-Nva-Ile-Arg-ProNHCH₂CH₃,
 (31) N-Ac-Sar-Gly-Val-D-Leu-Trp-Nva-Ile-Arg-ProNHCH₂CH₃,
 (32) N-Ac-Sar-Gly-Val-D-Leu-Tyr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (33) N-Ac-Sar-Gly-Val-D-Leu-Gly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (34) N-Ac-Sar-Gly-Val-D-Leu-2Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
 25 (35) N-Ac-Sar-Gly-Val-D-Leu-1Nal-Nva-Ile-Arg-ProNHCH₂CH₃,
 (36) N-Ac-Sar-Gly-Val-D-Leu-Octylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (37) N-Ac-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂CH₃,
 (38) N-Ac-Sar-Gly-Val-D-Leu-Allylgly-Nva-Ile-Arg-ProNHCH₂CH₃,
 (39) N-Ac-Sar-Gly-Val-D-Leu-D-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 30 (40) N-Ac-Sar-Gly-Val-D-Ile-Thr-Tyr-Ile-Arg-ProNHCH₂CH₃,
 (41) N-Ac-Sar-Gly-Val-D-Ile-Thr-Glu-Ile-Arg-ProNHCH₂CH₃,
 (42) N-Ac-Sar-Gly-Val-D-Ile-Thr-Propargylgly-Ile-Arg-ProNHCH₂CH₃,
 (43) N-Ac-Sar-Gly-Val-D-alloIle-Thr-Gln-Ile-Arg-ProNHCH₂CH₃,
 (44) N-Ac-Bala-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,

- (45) N-Phenylacetyl-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (46) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-AzaglyNH₂,
 (47) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-SerNH₂,
 (48) N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 5 (49) N-(6-Ac-Aca)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (50) N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (51) N-(4-Ac-Gaba)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (52) N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (53) N-(2-Furoyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 10 (54) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (55) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (56) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (57) N-(Shikimyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 15 (58) N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Gln-Ile-Arg-ProNHCH₂(CH₃)₂,
 (59) N-(2-Me-nicotinyl)-Sar-Gly-Val-D-Leu-Ser-Nva-Ile-Arg-ProNHCH₂(CH₃)₂,
 (60) N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-Pro-OH,
 (61) N-Ac-Sar-Ala-Val-D-Ile-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 20 (62) N-Ac-Sar-Gly-Val-D-Pen-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 (63) N-Ac-Sar-Gly-Val-D-Phe(3,4,5-triF)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃,
 and
 (64) N-Ac-Sar-Gly-Val-D-Phe(4-NH₂)-Thr-Nva-Ile-Arg-ProNHCH₂CH₃.

14. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

15. A method of treating a patient in need of anti-angiogenesis therapy comprising administering to the patient in need a therapeutically effective amount of a compound in Claim 1.


16. A composition for the treatment of a disease selected from cancer, arthritis, psoriasis, angiogenesis of the eye associated with infection or surgical intervention,

macular degeneration and diabetic retinopathy comprising a peptide as defined in Claim 1 in combination with a pharmaceutically acceptable carrier.

17. A method of isolating a receptor from an endothelial cell comprising binding a peptide as defined in Claim 1 to the receptor to form a peptide receptor complex; isolating the peptide receptor complex; and purifying the receptor.

Abstract

Peptides having the formula:



wherein A_0 is selected from hydrogen or an acyl group; A_{10} is a hydroxyl group or an amino acid amide; and A_1 , A_2 , A_3 , A_4 , A_5 , A_6 , A_7 , A_8 , and A_9 are amino acyl residues as defined herein.

**PATENT
IN THE UNITED STATES PATENT & TRADEMARK OFFICE**

APPLICANT: J. Henkin, et al. (Express Mail Label No. EL384166875US
)
SERIAL NO.: (not yet assigned) (I hereby certify that this paper (along with any
) paper referred to as being attached or enclosed) is
) being deposited with the United States Postal
FILED: November 22, 1999 (Service as Express Mail Post Office to Addressee
) Service under 37 C.F.R. 1.10 addressed to:
)
FOR: PEPTIDE ANTIANGIOGENIC (Box Patent Application
DRUGS) Assistant Commissioner for Patents
(Washington, D.C.
)
EXAMINER: (not yet assigned) ()
()
GROUP ART UNIT: (not yet assigned) (Date of Deposit: November 22, 1999
)
()
CASE NO.: 6356.US.P3 (Kathleen T. Litz 11/22/99
) Kathleen T. Litz Date
()
DATE: November 22, 1999 ()
()

**DECLARATION AND POWER OF ATTORNEY
FOR A UNITED STATES PATENT APPLICATION**

As a below-named inventor, I hereby declare:

My residence, post office address and citizenship are as stated below next to my name. I believe I am an original and first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention PEPTIDE ANTIANGIOGENIC DRUGS, the specification of which is attached.

I hereby state that I have reviewed and understand the contents of the above-mentioned specification, including the claims.

I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

Claim to benefit of foreign application(s) as follows:

I hereby claim foreign priority benefits under 35 U.S.C. §119 for the following foreign applications for patent or inventors certificate.

NONE

The following foreign applications for patent or inventors certificate have a filing date earlier than the filing date of the applications identified above.

NONE

Claim to benefit of earlier U.S. application(s) as follows:

I hereby claim the benefit under 35 U.S.C. §120 of the following earlier-filed United States patent applications. Insofar as the subject matter of each of the claims of this application is not disclosed in the prior U.S. applications in the manner required by 35 U.S.C. §112, first paragraph, I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which came into existence between the filing date(s) of the prior applications and the national or PCT filing date of this application.

Serial No. 09/316,888 filed May 21, 1999

I hereby appoint the following Attorneys and/or agents to prosecute this application and any continuation or divisional applications based hereon, and to transact all business in the Patent and Trademark Office connected therewith:

Mona Anand, Reg. No. 34,537
Regina M. Anderson, Reg. No. 35,820
Mark C. Bach, Reg. No. 34,766
Cheryl L. Becker, Reg. No. 35,441
Thomas D. Brainard, Reg. No. 32,459
Dianne Casuto, Reg. No. 40,943
Portia Chen, Reg. No. 44,075
Steven R. Crowley, Reg. No. 31,604
Andreas M. Danckers, Reg. No. 32,652
J. Michael Dixon, Reg. No. 32,410
Mimi C. Goller, Reg. No. 39,046
Daniel J. Hulseberg, Reg. No. 36,554

Neal D. Marcus, Reg. No. 35,267
James D. McNeil, Reg. No. 26,204
Robert A. Miller, Reg. No. 26,956
Lawrence S. Pope, Reg. No. 26,791
Nicholas A. Poulos, Reg. No. 30,209
Dugal S. Sickert, Reg. No. 33,784
Gregory W. Steele, Reg. No. 33,796
Michael J. Ward, Reg. No. 37,960
David L. Weinstein, Reg. No. 28,128
Steven F. Weinstock, Reg. No. 30,117
Brian R. Woodworth, Reg. No. 33,137
Paul D. Yasger, Reg. No. 37,477

Send Correspondence to:

Steven F. Weinstock
D-377/AP6D
Abbott Laboratories
100 Abbott Park Road
Abbott Park, IL 60064-6050

Direct telephone calls to:

Portia Chen
(847) 937-4558

Name: Jack Henkin
Post Office Address: 1370 Lincoln Avenue South, Highland Park, IL 60035
Residence: Highland Park, IL 60035
Citizenship: United States of America

Name: Fortuna Haviv
Post Office Address: 1125 Oxford Road, Deerfield, Illinois 60015
Residence: Deerfield, Illinois 60015
Citizenship: United States of America

Name: Michael F. Bradley
Post Office Address: 3930 N. Pine Grove Ave., Apt. #2405, Chicago, IL 60613
Residence: Chicago, Illinois 60613
Citizenship: United States of America

Name: Douglas M. Kalvin
Post Office Address: 1201 Lockwood Drive, Buffalo Grove, Illinois 60089
Residence: Buffalo Grove, Illinois 60089
Citizenship: United States of America

Name: Andrew J. Schneider
Post Office Address: 980 Cheswick Drive, Gurnee, Illinois 60031
Residence: Gurnee, Illinois 60031
Citizenship: United States of America

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that all statements made herein were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Jack Henkin

Date

Fortuna Haviv

Date

Michael F. Bradley

Date

Douglas M. Kalvin

Date

Andrew J. Schneider

Date